

J. C. MIRALLES LÓPEZ, A. I. ESCUDERO PASTOR, A. CARBONELL MARTÍNEZ,  
C. NAVARRO GARRIDO, Y. BONILLA PACHECO, Y. PETRYK PETRYK

# Quality of life in severe asthmatic patients treated with benralizumab

Allergy Section, University General Hospital Reina Sofia, Murcia, Spain

## KEY WORDS

*Quality of life; severe asthma; benralizumab; control correlation; lung function correlation.*

## Corresponding author

Juan Carlos Miralles López  
Allergy Section  
University General Hospital Reina Sofia  
Avenida Intendente Jorge Palacios 1  
30003 Murcia, Spain  
ORCID ID: 0000-0001-8811-3939  
E-mail: jucarmir@telefonica.net

## Doi

10.23822/EurAnnACI.1764-1489.218

To the Editor,

severe asthma has a great impact on the health related quality of life (HRQoL) of patients and their families (1). Thus, an appropriate asthma control is expected to improve HRQoL. Biologics, primarily monoclonal antibodies, have been developed to target specific pathways and important molecules in the pathogenesis of asthma, and their use has shown some promising effects on the HRQoL of severe asthmatic patients (2).

Benralizumab treatment decreases exacerbations and improves lung function in patients with severe, uncontrolled eosinophilic asthma (3), and we have reported in a real-life study of 10 patients with severe asthma (4), that benralizumab decreases the number of exacerbations improving disease control.

The aim of this work is to study the effect of benralizumab in the HRQoL of patients with severe asthma, its different domains, and the relationship of the improvement in HRQoL with the changes in other variables such as asthma control, exacerbations, and lung function. The study was conducted in routine clinical practice, in accordance with the ethical standards established in the Declaration of Helsinki of 1946, and informed consent was obtained from all participants before enrolment in the study.

We have measured epidemiological variables, eosinophilia, HRQoL, number of exacerbations, asthma control (Asthma

Control Test questionnaire), and lung function. We have assessed HRQoL by applying the mini AQLQ (Asthma Quality of Life Questionnaire); this questionnaire covers four health domains: symptoms, activity limitation, emotional function, and discomfort due to environmental stimuli. The nonparametric Wilcoxon signed-rank test has been used for the statistical analysis and the results are described by median and interquartile range (IQR). We have correlated the changes in quality of life data with the changes in other variables such as lung function, asthma control and exacerbations using the Spearman's rank correlation coefficient. We present data from 15 patients (9 women, 6 men) with severe eosinophilic bronchial asthma who have received treatment with benralizumab for 6 months. The mean age was 58.8 years (range 39-78), nine were never smokers and 6 ex-smokers. Most of them had overweight with an average BMI of 33.14.

Regarding background, 10 patients referred chronic rhinosinusitis, 8 nasal polyposis, and eight were atopic. About "asthma severity", nine patients were in step 5 of GEMA – Spanish Guide for Asthma Management – (treated with high-dose inhaled corticosteroids and long-term bronchodilators, in addition to antileukotrienes and anticholinergics), and other six patients were in step 6, requiring continuous oral steroids.

Referring to adverse effects, administration of benralizumab was uneventful for 14 patients, but one developed mild fever

controlled with paracetamol. This good tolerance agrees with previous reports on the safety of the drug (5).

Mean eosinophil blood count was 522 cells/ $\mu$ L before the drug and 48 cells/ $\mu$ L after it, with 9 patients presenting 0 eosinophils. The results of the mini AQLQ are shown in the **table I**. We have obtained a statistically significant improvement in the total score: median pre-treatment 2.93 and post-treatment 5.60 ( $p < 0.001$ ), and in the four domains: "symptoms" 3.00 and 6.20 ( $p < 0.001$ ), "activity limitation" 3.00 and 5.75 ( $p < 0.001$ ), "emotional function" 2.67 and 6.00 ( $p = 0.001$ ), and "environmental stimuli" 3.33 and 5.67 ( $p = 0.043$ ).

All the patients achieved the minimal important difference (6) of improvement in the AQLQ Total score, 14 patients in the "limitation activity" domain, 13 patients in the "symptoms" domain, 12 in the "emotional function" domain and 8 patients in the "environmental stimuli" domain.


All patients experienced clinical improvement with the treatment. The number of exacerbations decreased in all cases, and the median for the year prior to benralizumab was 3 (2.00-4.00) and 0 (0.00-0.00) (annualized rate) after the drug ( $p = 0.001$ ).

We also obtained improvements in asthma control: median ACT pre-treatment 12 (8.00-16.00), post-treatment 21 (19.00-23.00) ( $p < 0.001$ ) and non-significant for lung function: median FEV1 pre-treatment 1.6 L (1.14-2.06), post-treatment 1.81 L (1.19-2.93) ( $p = 0.078$  NS). We have correlated the changes in the Total AQLQ score and the four domains with the changes in asthma control, lung function and exacerbations. The total AQLQ score shows significant correlation with ACT:  $r_s = 0.615$  ( $p = 0.015$ ) and FEV1:  $r_s = 0.739$  ( $p = 0.002$ ), but not with decrease of exacerbations:  $r_s = -0.430$  ( $p = 0.109$  NS) (**figure 1**). Similar results have been obtained by "symptoms", which also correlates with ACT:  $r_s = 0.535$  ( $p = 0.040$ ) and FEV1:  $r_s = 0.702$  ( $p = 0.004$ ), but not with exacerbations:  $r_s = -0.419$  ( $p = 0.120$  NS), and "environmental stimuli" with ACT:  $r_s = 0.534$  ( $p = 0.040$ ), with FEV1:  $r_s = 0.591$  ( $p = 0.020$ ), and exacerbations:  $r_s = -0.349$  ( $p = 0.203$  NS). Nevertheless, the domain "activity limitation" correlates with the three variables, ACT:  $r_s = 0.706$  ( $p = 0.003$ ), FEV1:  $r_s = 0.680$  ( $p = 0.005$ ), and exacerbations  $r_s = -0.518$  ( $p = 0.048$ ). Finally, the domain "emotional function" does not correlate with any other variable: ACT:  $r_s = 0.325$  ( $p = 0.238$  NS), FEV1:  $r_s = 0.504$  ( $p = 0.055$  NS), and exacerbations  $r_s = -0.286$  ( $p = 0.302$  NS).

**Table I** - Results of mini AQLQ questionnaire.

AQLQ	Pre	Post	DIF	P-value
<b>Symptoms</b>	3.00 (2.20-4.20)	6.20 (5.80-6.80)	3.20	< 0.001
Shortness of breath (1)	3	6		
Discomfort due to coughing (4)	2	6		
Chest tightness (6)	4	7		
Sleep disturbance (8)	3	7		
Wheezes (10)	2	6		
<b>Activity Limitation</b>	3.00 (1.25-4.00)	5.75 (4.50-6.75)	2.75	0.001
Intense efforts (12)	1	5		
Moderate efforts (13)	3	6		
Social activities (14)	4	7		
Work activities (15)	3	6		
<b>Emotional Function</b>	2.67 (1.67-3.67)	6.00 (4.33-7.00)	3.33	0.001
Frustrated-irritated (3)	3	6		
Fear not having medication (5)	2	6		
Concerned about asthma (9)	2	6		
<b>Environmental Stimuli</b>	3.33 (2.67-6.33)	5.67 (4.00-6.67)	2.34	0.043
Discomfort due to Dust (2)	6	6		
Discomfort due to Tobacco (7)	4	7		
Discomfort due to Pollution (11)	3	6		
<b>Total</b>	2.93 (2.67-4.27)	5.60 (4.73-6.87)	2.67	< 0.001

Median (Interquartile range); question number in questionnaire in parenthesis.

**Figure 1** - Correlation between AQLQ and ACT, FEV1 and Exacerbations.

We have obtained, after treatment with benralizumab, a significant improvement in the patients' HRQoL, both Total AQLQ score and the four domains "symptoms", "activity limitation", "emotional function" and "environmental stimuli". This improvement correlates with improvement in asthma control (ACT) and lung function in the case of total score, "symptoms" and "environmental stimuli", and in addition with decrease of exacerbations in "activity limitation".

In accordance with other authors, a correlation was found between improvement in HRQoL and asthma control (7) and lung function improvement. The lack of correlation of the changes in HRQoL with decrease of exacerbations (but "activity limitation") has also been reported by Enríquez-Matas *et al.* (8), and it could be due to the fact that the short period covered for the AQLQ questionnaire (two weeks) makes more difficult reflecting changes in a variable such as exacerbations. This limitation is more evident considering the long free period there is sometimes between exacerbations.

We conclude that benralizumab significantly improves HRQoL in severe asthmatic patients, globally and by domains: "symptoms", "activity limitation", "emotional function", and discomfort due to "environmental stimuli". This improvement is correlated with better asthma control and lung function.

### Fundings

None.

### Conflict of interests

Juan Carlos Miralles López has received lecture fees from Novartis, GSK, Astra Zeneca, Sanofi and Chiesi. The rest of authors declare that they have no conflict of interests.

### References

1. Stucky BD, Sherbourne CD, Edelen MO, Eberhart NK. Understanding asthma-specific quality of life: moving beyond asthma symptoms and severity. *Eur Respir J.* 2015;46(3):680-7. doi: 10.1183/09031936.00225014.
2. Hossny E, Carballo L, Casale T, El-Gamal Y, Rosenwasser L. Severe asthma and quality of life. *World Allergy Organ J.* 2017;10(1):28. doi: 10.1186/s40413-017-0159-y.
3. Chippes BE, Newbold P, Hirsch I, Trudo F, Goldman M. Benralizumab efficacy by atopy status and serum immunoglobulin E for patients with severe, uncontrolled asthma. *Ann Allergy Asthma Immunol.* 2018;120(5):504-511.e4. doi: 10.1016/j.anai.2018.01.030.
4. Miralles López JC, Escudero Pastor AI, Carbonell Martínez A, Navarro Garrido C, Bonilla Pacheco Y, Petrik Petrik Y. Benralizumab in Real Life. *J Investig Allergol Clin Immunol.* 2021;31(1):87-88. doi: 10.18176/jiaci.0599.
5. Busse WW, Bleecker ER, FitzGerald JM, Ferguson GT, Barker P, Sproule S, *et al*; BORA study investigators. Long-term safety and efficacy of benralizumab in patients with severe, uncontrolled asthma: 1-year results from the BORA phase 3 extension trial. *Lancet Respir Med.* 2019;7(1):46-59. doi: 10.1016/S2213-2600(18)30406-5.
6. Juniper EF, Guyatt GH, Willan A, Griffith LE. Determining a minimal important change in a disease-specific Quality of Life Questionnaire. *J Clin Epidemiol.* 1994;47(1):81-7. doi: 10.1016/0895-4356(94)90036-1.
7. Howell CR, Thompson LA, Gross HE, Reeve BB, Huang SW, DeWalt DA, *et al.* Association of consistently suboptimal quality of life with consistently poor asthma control in children with asthma. *Ann Allergy Asthma Immunol.* 2017;119(6):562-564.e1. doi: 10.1016/j.anai.2017.09.053.
8. Enríquez-Matas A, Fernández-Rodríguez C, Andrés Esteban EM, Fernández-Crespo J. Main Contributory Factors on Asthma Control and Health-Related Quality of Life in Elderly Asthmatics. *J Investig Allergol Clin Immunol.* 2020;30(4):264-271. doi: 10.18176/jiaci.0430.