

4. Sonkoly E, Muller A, Lauerma AI, et al. IL-31: a new link between T cells and pruritus in atopic skin inflammation. *J Allergy Clin Immunol.* 2006;117:411-7.
5. Raap U, Wieczorek D, Gehring M, et al. Increased levels of serum IL-31 in chronic spontaneous urticaria. *Exp Dermatol.* 2010;19:464-6.
6. Rabenhorst A, Hartmann K. Interleukin-31: a novel diagnostic marker of allergic diseases. *Curr Allergy Asthma Rep.* 2014;14:423.
7. Gangemi S, Franchina T, Minciullo PL, et al. IL-33/IL-31 axis: a new pathological mechanisms for EGFR tyrosine kinase inhibitors-associated skin toxicity. *J Cell Biochem.* 2013;114:2673-6.
8. Taniguchi K, Yamamoto S, Hitomi E, et al. Interleukin 33 is induced by tumor necrosis factor alpha and interferon gamma in keratinocytes and contributes to allergic contact dermatitis. *J Investig Allergol Clin Immunol.* 2013;23:428-34.
9. Mattii M, Ayala F, Balato N, et al. The balance between pro- and anti-inflammatory cytokines is crucial in human allergic contact dermatitis pathogenesis: the role of IL-1 family members. *Exp Dermatol.* 2013;22:813-9.
10. Gangemi S, Merendino RA, Minciullo PL, et al. Serum levels of interleukin-18 in subjects affected by occupational allergic contact dermatitis. *J Dermatol Sci.* 2003;33:187.
11. Ricciardi L, Minciullo PL, Saitta S, et al. Increased serum levels of IL-22 in patients with nickel contact dermatitis. *Contact Dermatitis.* 2009;60:57-8.
12. Stott B, Lavender P, Lehmann S, et al. Human IL-31 is induced by IL-4 and promotes TH2-driven inflammation. *J Allergy Clin Immunol.* 2013;132:446-54.
13. Komai-Koma M, Gilchrist DS, McKenzie ANJ et al. IL-33 activates B1 cells and exacerbates contact sensitivity. *J Immunol.* 2011;186:2584-91.