

ABSTRACT

Atopic dermatitis is a common illness in pediatric age. Children with atopic dermatitis are prone to develop cutaneous sensitization due to skin barrier dysfunction and immune dysregulation. Recent studies have shown a higher prevalence of certain allergens, which identification may be clinically relevant and have implications for atopic dermatitis management. Considering the most prevalent and relevant allergens based on a retrospective analysis of 145 pediatric patients, 44,1% (n=63) with atopic dermatitis, and comparing the positive results, we propose the application of an adapted baseline series with the most relevant 20 allergens for the Portuguese pediatric population with atopic dermatitis with recommendation for an evaluation of allergic contact dermatitis.

KEYWORDS

Atopic dermatitis, allergic contact dermatitis, prevalent allergens, baseline serie, pediatric patients.

IMPACT STATEMENT

Children with and without atopic dermatitis have similar contact sensitization rates however to different allergens. An adapted baseline series with the most relevant allergens is proposed for the pediatric population with atopic dermatitis.

LIST OF ABBREVIATIONS

AD - Atopic dermatitis

ACD - Allergic contact dermatitis

EAACI - European Academy of Allergy and Clinical Immunology

ESCD - European Society of Cutaneous Allergy and Contact Dermatitis

ICDRG - International Contact Dermatitis Research Group

INTRODUCTION

Atopic dermatitis (AD) is a common illness in pediatric population. In contrast, allergic contact dermatitis (ACD) used to be considered a rarity (1-3). Children with AD are prone to develop cutaneous sensitization due to several factors, due to skin barrier disruption, immune dysregulation, and consequent increased penetration of allergens (1-3). Children with AD are exposed to several allergens from an early age. The frequent application of numerous topical treatments such as emollients and topical drugs may determine a characteristic sensitization profile (1,3,4).

Recent studies have shown that sensitization rates are similar between children with and without AD, although certain allergens have a higher prevalence in AD patients. Despite this, the interrelation of AD and ACD is complex and controversy remains over whether AD increases the risk of contact allergy in children or determines different contact sensitization profiles (1,2).

Epicutaneous patch testing is the gold standard test for the diagnosis of ACD. It is a specific, relatively inexpensive and safe procedure, even for preschool children however, one of the main limitations is the limited body surface area available for adhesive application (4). Therefore, a selection of allergens based on the exposures and clinical examination are recommended but difficult to standardize (1,4).

The identification of clinically relevant allergen sensitization in a patient with AD may have important implications for AD disease management and symptom control (3-5). According to the European Academy of Allergy and Clinical Immunology (EAACI) position paper published in 2015 and other recent reviews (1,4,6), epicutaneous patch testing is recommended in patients with AD with:

- a suspicion of specific ACD;
- difficult-to-control AD;
- new-onset dermatitis in patients with AD;
- prior to initiating systemic therapy for severe AD;

Considering the difficulty in identifying relevant allergens and their standardization, several research groups have proposed a “baseline series” for the pediatric population, based on sensitization rates in Europe and specific regions (4,7,8). Regarding the Portuguese population, few data is available (1,3), therefore the aim of this study was the creation and subsequent application of a “baseline series” for pediatric AD patients

with suspected ACD, difficult-to-control AD, new-onset dermatitis and prior to initiating systemic therapy for AD, or when no specific exposure is identified, in order to improve AD management and treatment in pediatric patients.

MATERIALS AND METHODS

Study design

The authors performed a retrospective, descriptive and inferential review of pediatric patients (<18 years old) patch tested at the Contact Allergology Unit of the Dermatology and Venereology Department in a tertiary hospital, between 2005 and 2021 (17 years). As a result of the evaluation of our data, we propose a baseline series for evaluation in pediatric AD patients.

All legal guardians of pediatric patients received written informed consent. This paper was written considering the ethical and legal principles and following the recommendations of the Declaration of Helsinki of the World Medical Association. The anonymity of all the participants in this work was guaranteed.

Subjects

Patients were characterized according to demographic data, personal and family history of atopy, and clinical and diagnostic parameters such as allergen series applied, positive reactions, and relevance. Atopic dermatitis was diagnosed according to the criteria of Hanifin and Rajka (9).

All patients included were routinely tested with the European baseline series and/or with the cosmetic series. The allergens used were from Chemotechnique Diagnostics[®], Vellinge, Suécia; Trolab Allergens[®], Smartpractice GmbH, Alemanha; Bial Alergénios[®]. Although, each series has been selected and adapted following the European Society of Cutaneous Allergy and Contact Dermatitis (ESCD) recommendations. Additionally, and in accordance with the available body surface and clinical history, supplementary series or specific allergens were applied. Allergen chambers were from (Epitest Ld[®]) or IQUltra chambers (Chemotechnique diagnostics[®]). The allergen chambers were applied on intact upper back and kept in occlusion for 48 hours. Readings were carried out on day(D) 3 or D4, and, in specific cases, associated with a second reading on D7. The

reactions were scored according to the recommendations of the International Contact Dermatitis Research Group (ICDRG) and ESCD (10).

It was assessed the sensitization prevalence rates for the population as a whole and stratified for children with and without AD.

Allergens patch tested <20 times were excluded from this analysis to avoid non-inferential results. A value of 20 was chosen to reach a comparable number of patients tested with at least substances in the European baseline series.

Statistical analysis

Statistical analysis was performed using SPSS Statistics version 24.0®. Descriptive statistics were analysed as measures of central tendency and dispersion. The comparison of proportions was made using the chi-square test. Fisher's exact test was used in cases of low sample sizes. A Type I error of 0.05 was considered.

The study population was selected and compared into two groups: patients with AD and patients without AD. Patients included in the AD group met the Hanifin and Rajka criteria, with current atopic dermatitis or a history of atopic dermatitis (9). The clinical relevance of patch tests results was determined for each positive patch test result, according to the clinical history, and considering if there was a suspicion that the allergen would exacerbate or trigger the presence of dermatitis.

RESULTS

Demographic and clinical characteristics

In a total sample of n=145 pediatric patients, 68.3% (n=99) females, with a median 13.0 years old (Interquartile range 10-15), corresponding to 4,1% (n=6) with < 6 years, 35,2% (n=11) with 6-12 years, and 60,7% (n=88) with >12 years, were included in the analysis. AD patients corresponding to 43.4% (n=63) of the total sample. Patient characteristics are summarized in Table 1.

Major location of suspected ACD

Regarding the affection of suspected ACD body areas (table 1), the face corresponded to the main suspected body area in both groups. Particularly, the body area corresponding to the hands was more representative in the group of patients with AD;

AD group 17.5% (n=11) vs. non-AD group 36.6% (n=30), $p < 0.001$. Suspicion of ACD associated with the periumbilical location has only been described in patients with AD; AD group 6.3% (n=4) versus non-AD group 0.0% (n=0), $p < 0.046$.

Patch-testing results

The presence of allergic sensitization was similar in both groups. At least the presence of one allergic sensitization was demonstrated in 48.3 % (n=70), corresponding to 49.2% (n=31) to AD patients (Table 1).

In table 2 are shown the allergens tested and their sensitization prevalence rates for the total sample, AD patients group and non-AD patients group. In table 3 are listed the 20 most common sensitization prevalence rates for each group. The most prevalent allergen for both groups was nickel sulfate. Between the two groups evaluated, there were variable prevalences among the different allergens tested.

In the AD group there was a higher prevalence of sensitization to: quaternium 15, 1% petrolatum, parthenolide, diazolidinyl urea, 2-hydroxyethyl methacrylate, hydroperoxide of limonene, compositae mix II, 1,2-dibromo-2,4-dicyanobutane, caine mix II, sesquiterpene lactone mix, and epoxy resin, sensitizations that were not prevalent in the non-AD group. In AD group, none of the most prevalent allergens were included in cosmetic series.

In contrast, allergens such as hydroperoxide linalool, methylchloroisothiazolinone/methylisothiazolinone, fragrance mix I, amercchol I-101, disperse orange 3, p-phenylenediamine, imidazolidinyl urea, formaldehyde, and n-isopropyl-n'-phenyl-phenylenediamine had a similarly high prevalence in both analysed groups.

Considering the clinical relevance, based on the criteria mentioned above, table 4 shows the 20 main relevant allergens for each group separately. The most prevalent allergen, nickel sulfate, was considered among the 20 most relevant tested allergens however, it assumed a minor position due to its high prevalence.

The tested allergens methylchloroisothiazolinone/methylisothiazolinone, fragrance mix I, diazolidinyl urea, p-Phenylenediamine, imidazolidinyl urea, methylisothiazolinone, nickel sulfate, and hydroperoxide linalool proved to be relevant in both groups studied. In the AD group, the tested allergens quaternium 15, parthenolide, 2-Hydroxyethyl

methacrylate, hydroperoxide of limonene, compositae mix II, 1,2-Dibromo-2,4-dicyanobutane, caine mix II, N-Isopropyl-N'-phenyl-phenylenediamine, sesquiterpene lactone mix, potassium dichromate, parabens, and amerchol L-101 differ from the non-AD group. Based on the results presented, Figure 1 shows the new proposed baseline series for AD patients with a recommendation for ACD evaluation.

DISCUSSION

In this study, we report the experience of our hospital in patch test evaluation in the pediatric population. In our study, most patch-tested patients were aged between 10 and 15 years old. The AD group included significantly younger patients, with most patients aged between 9 and 14 years old. These findings support the notion that ACD increases with age in general and that ACD affects AD patients at an earlier age. They are in line with the probability that AD patients, whose disease presentation appears at a younger age, may have an increased risk of ACD (1,3). However, the percentage of positive tests in different age groups was similar. In addition, although the safety of patch testing has been verified in children from 6 months of age, the reduced number of children tested at ages below 6 years may reflect the lack of knowledge of the safety profile in this age group (10). Therefore, it is important to the awareness of health professionals for early recognition of this clinical presentation.

Despite the interrelation between AD and ACD is controversial, the failure to identify a potential aggressor allergen can have important consequences for the evolution and management of AD. Some of the most common allergens shown in our sample, such as nickel, cobalt, methylchloroisothiazolinone/methylisothiazolinone, hydroperoxide linalool, fragrances, p-phenylenediamine, quaternium 15 and formaldehyde, have been similarly described in the literature as having a high prevalence (11). Regarding patients with and without atopic dermatitis, the contact sensitization rates were similar however, after nickel, main allergens were different, mostly related to preservatives (formaldehyde releasers) and plant extracts (parthenolide) that may be present in topical products used in the management of AD. Pointing out particular allergens, nickel as the most prevalent allergen for both groups, is a known commonly identified allergen in many studies of ACD in children and adolescents, likely attributable to the increased use of jewellery, children's toys, metallic components of children's clothing, electronic

equipment, and dental appliances (12). Sensitizations to methylisothiazolinone and cobalt were more prevalent in the non-AD group. Although methylchloroisothiazolinone/methylisothiazolinone was one of the most prevalent in both groups, the individual sensitization to methylisothiazolinone showed to be more frequent in non-AD group, which is described in previous studies as a frequent allergen present in ACD of hands and face due to its activity against bacteria and fungi (13). Regarding the prevalence of sensitization to cobalt, our data diverges from some studies published (11,14). For example, in China it was found that sensitization to cobalt was higher in patients with AD and was dependent on geographic exposures, not being necessarily related to the coexistence of sensitization to other metals such as nickel (14). Our study has some limitations that are considered transversal to the other works presented on this topic, namely the possibility of differences in reading techniques between observers. However, in our work, one of the members of the evaluation of the results will have always been the same to limit this bias. Another limitation may be the difficulty distinguishing between allergic or irritant reactions. The data collected in this work involved children from the same region and a considerable period of time, which may influence the results. We need to consider the high variability of allergen expression in the environment, changes in European legislation, modifications in components of topical products, and lifestyle habits.

CONCLUSION

Confirmed allergic contact dermatitis was relatively common in our sample. It appeared to increase with age however, the rates of positive tests were similar between age groups. As expected, there were notable differences in sensitization rates of relevant allergens when compared to other European studies, confirming the geographic sensitization variability and need for pediatric adaptation and custom “baseline series”. Patients with and without atopic dermatitis had similar contact sensitization rates however the main allergens were different. We propose the first Portuguese Pediatric Baseline Series adapted for AD patients, based on the most common and relevant allergens. Overall, this study provides key information on the relationship between AD and ACD in pediatric patch-tested patients in Portugal. Further studies on a national

scale are necessary to elucidate and validate this new serie proposal for the Portuguese pediatric population.

REFERENCES

1. Simonsen AB, Johansen JD, Deleuran M, Mortz CG, Sommerlund M. Contact allergy in children with atopic dermatitis: a systematic review. *Br J Dermatol*. 2017;177(2):395-405. doi:10.1111/bjd.15628
2. Rodrigues DF, Goulart EM. Patch-test results in children and adolescents: systematic review of a 15-year period. *An Bras Dermatol*. 2016;91(1):64-72. doi:10.1590/abd1806-4841.20163927
3. Borok J, Matiz C, Goldenberg A, Jacob SE. Contact Dermatitis in Atopic Dermatitis Children-Past, Present, and Future. *Clin Rev Allergy Immunol*. 2019;56(1):86-98. doi:10.1007/s12016-018-8711-2
4. De Waard-van der Spek FB, Darsow U, Mortz CG, Orton D, Worm M, Muraro A, et al. EAACI position paper for practical patch testing in allergic contact dermatitis in children. *Pediatr Allergy Immunol*. 2015; 26(7): 598-606. doi:10.1111/pai.12463
5. Steuer MS, Botto NC. Patient Reported Improvement After Patch Testing and Allergen Avoidance Counseling: A Retrospective Analysis. *Dermatol Ther (Heidelb)*. 2018. 8: 435–440. doi:10.1007/s13555-018-0250-5
6. Hamann CR, Hamann D, Egeberg A, Johansen JD, Silverberg J, Thyssen JP. Association between atopic dermatitis and contact sensitization: a systematic review and meta-analysis. *J Am Acad Dermatol*. 2017;77(1):70–8. doi:10.1016/j.jaad.2017.02.001

7. Belloni Fortina A, Cooper SM, Spiewak R, Fontana E, Schnuch A, Uter W. Patch test results in children and adolescents across Europe. Analysis of the ESSCA Network 2002-2010. *Pediatr Allergy Immunol.* 2015; 26(5):446-55. doi:10.1111/pai.12397
8. Jacob SE, Lipp MB, Suh E, Goldenberg A. Practice Patterns of Dermatologists in the Pediatric Contact Dermatitis Registry. *Pediatr Dermatol.* 2017;34(4):408-12. doi:10.1111/pde.13154
9. Akan A, Dibek-Mısırlıoğlu E, Civelek E, Vezir E, Kocabaş CN. Diagnosis of atopic dermatitis in children: comparison of the Hanifin-Rajka and the United Kingdom Working Party criteria. *Allergol Immunopathol.* 2020; 48(2):175-181. doi:10.1016/j.aller.2019.07.008
10. Johansen DJ, Aalto-Korte K, Agner T, Andersen KE, Bircher A, Bruze M, et al. European Society of Contact Dermatitis guideline for diagnostic patch testing – recommendations on best practice. *Contact Dermatitis.* 2015; 73:195-221. doi:10.1111/cod.12432
11. Admani S, Jacob SE. Allergic contact dermatitis in children: review of the past decade. *Curr Allergy Asthma Rep.* 2014; 14(4):421. doi:10.1007/s11882-014-0421-0
12. Jacob SE, Goldenberg A, Pelletier JL, Fonacier LS, Usatine R, Silverberg N. Nickel Allergy and Our Children's Health: A Review of Indexed Cases and a View of Future Prevention. *Pediatr Dermatol.* 2015; 32(6):779-85. doi:10.1111/pde.12639
13. Schwensen JF, Uter W, Bruze M, Svedman C, Goossens A, Wilkinson M. The epidemic of methylisothiazolinone: a European prospective study. *Contact Dermatitis.* 2017; 76:272–9. doi:10.1111/cod.12733
14. Peng Fen S, Robert A, Chen Z, Zhang JZ. High prevalence of contact hypersensitivity to metals and preservatives in Chinese patients with atopic dermatitis, *Chinese Medical Journal.* 2019; 132(23):2881-2882. doi:10.1097/CM9.0000000000000526

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

FUNDING

No funding sources to declare.

Manuscript accepted for publication

Table 1. Demographics of Paediatric patients referred for Patch Testing due to ACD suspicion

Characteristic	Total	With atopic dermatitis	Without atopic dermatitis	p-value
N, % (n)	100.0 (145)	43.4 (63)	56.6 (82)	
Sex (female), % (n)	68.3 (99)	65.1 (41)	70.7 (58)	0.472
Age (years), median (IQR)	13.0 (10.0-15.0)	12.0 (9.0-14.0)	14.5 (11.0-16.3)	0.015
Group age				
< 6 years, % (n)	4.1 (6)	4.8 (3)	3.7 (3)	>0.99
6-12 years, % (n)	35.2 (51)	44.4 (28)	45.1 (23)	0.467
>12 years, % (n)	60.7 (88)	50.8 (32)	63.6 (56)	0.104
Asthma, % (n)	25.5 (37)	28.6 (18)	23.2 (19)	0.463
Rhinitis, % (n)	25.5 (37)	38.1 (24)	15.9 (13)	0.02
Atopy, % (n)	50.3 (73)	76.2 (48)	30.5 (25)	0.001
Atopy family history, % (n)	31.7 (46)	42.9 (27)	23.2 (19)	0.011
Major location of suspected ACD				
Face, % (n)	40.0 (58)	42.9 (27)	37.8 (31)	0.232
Hands, % (n)	28.2 (41)	17.5 (11)	36.6 (30)	0.001
Upper or lower extremities, % (n)	13.1 (19)	12.7 (8)	13.4 (11)	0.648
Generalized, % (n)	6.8 (10)	9.5 (6)	4.9 (4)	>0.99
Flexures, % (n)	6.2 (9)	7.9 (5)	4.9 (4)	>0.99
Torso, % (n)	2.8 (4)	3.2 (2)	2.4 (2)	>0.99
Periumbilical, % (n)	2.8 (4)	6.3 (4)	0 (0)	0.046
≥1 positive patch test result, % (n)	48.3 (70)	49.2 (31)	47.6 (39)	0.868
≥ 2 positive patch test result, % (n)	20.0 (29)	17.5 (11)	21.2 (18)	0.537
≥ 3 positive patch test result, % (n)	11.7 (17)	7.9 (5)	14.6 (12)	0.299

P-value is result of differences between patients with and without atopic dermatitis groups.
Statistically significant values were marked bold.

Table 2. Overview of all routinely tested allergens (European baseline series and cosmetic series), and distribution of positive patch test reactions in the total paediatric population and in with and without atopic dermatitis.

Allergen	Total (n = 145)	With atopic dermatitis (n = 63)		Without atopic dermatitis (n = 82)		p-value
	% (n positive/ n tested)	% (n positive/ n tested)	% PRPP	% (n positive/ n tested)	% PRPP	
Baseline series						
Nickel sulfate 5% pet	16.0 (23/144)	16.1 (10/62)	70	15.9 (13/82)	62	>0.99
MCI/MI 0.02% aq	8.3 (12/144)	6.5 (4/62)	100	9.8 (8/82)	88	0.578
Hydroperoxide linalool 1/0,5% pet	6.8 (3/44)	7.4 (2/27)	50	5.6 (1/18)	100	>0.99
Methylisothiazolinone 0.2% aq	6.3 (9/144)	1.6 (1/63)	100	9.9 (8/81)	88	0.046
Amerchol L-101 50% pet	4.9 (7/143)	4.8 (3/62)	67	4.9 (4/81)	25	>0.99
Cobalt chloride 1% pet	4.9 (7/142)	0.1 (4/62)	50	3.8 (3/80)	33	0.699
Benzisotiazolinona 0.1% pet	4.9 (2/41)	0 (0/21)	NA	10.0 (2/20)	100	0.232
Fragance mix I 8% pet	4.8 (7/145)	4.8 (3/63)	100	4.9 (4/82)	75	>0.99
Caine mix III 10% pet	4.3 (6/141)	0 (0/60)	0	7.4 (6/81)	17	0.038
p-Phenylenediamine 1% pet	4.3 (6/140)	3.3 (2/61)	100	5.1 (4/79)	100	0.697
Disperse Orange 3 1% pet	3.8 (4/106)	4.8 (2/42)	50	3.1 (2/64)	0	0.648
Formaldehyde 2% aq	3.5 (5/143)	3.2 (2/62)	50	3.7 (3/81)	67	>0.99
Parthenolide 0.1% pet	3.4 (1/29)	6.3 (1/16)	100	0 (0/13)	NA	>0.99
Diazolidinyl urea 2% pet	2.8 (4/144)	4.8 (3/63)	100	1.2 (1/81)	100	0.319
Quaternium 15 1% pet	2.8 (4/141)	6.7 (4/60)	100	0 (0/81)	NA	0.031
Sodium disulfite 1% pet	2.8 (1/36)	0 (0/18)	NA	5.6 (1/18)	0	>0.99
N-Isopropyl-N'-phenyl-pphenylenediamine 0.1% pet	2.2 (3/138)	1.7 (1/59)	100	2.5 (2/79)	0	>0.99
Hidroperoxide of limonene 0.2/0.3% pet	2.2 (1/46)	3.7 (1/27)	100	0 (0/19)	NA	>0.99
Lanolin alcohols 30% pet	2.1 (3/144)	1.6 (1/63)	0	2.5 (2/81)	50	>0.99
Fragance mix II 14% pet	2.1 (3/144)	0 (0/63)	NA	3.7 (3/81)	67	0.257
Imidazolidinyl urea 2% pet	2.1 (3/141)	3.3 (2/61)	100	1.3 (1/80)	100	0.578
2-Hydroxyethyl methacrylate 1% pet	2.0 (1/49)	4.2 (1/24)	100	0 (0/25)	NA	0.49
Peru balsam 25% pet	1.4 (2/143)	1.6 (1/62)	0	1.2 (1/81)	100	>0.99
Compositae mix II 2.5% pet	1.1 (1/87)	2.6 (1/38)	100	0 (0/49)		0.437
Hydroxyisohexyl 3-cyclohexene carboxaldehyde (lyral) 5% pet	1.0 (1/98)	0 (0/47)	NA	1.9 (1/51)	100	>0.99
Propolis 10% pet	0.9 (1/109)	0 (0/46)	NA	1.6 (1/63)	100	>0.99
1,2-Dibromo-2,4-dicyanobutane 0,3% pet	0.8 (1/129)	1.8 (1/55)	100	0 (0/74)	NA	0.426
Caine mix II 10% pet	0.8 (1/128)	1.8 (1/55)	100	0 (0/73)	NA	0.43
Parabens 12% pet	0.7 (1/142)	1.6 (1/61)	100	0 (0/81)	NA	0.43
Potassium dichromate 0.5% pet	0.7 (1/142)	1.6 (1/62)	100	0 (0/80)	NA	0.437
Sesquiterpene lactone mix 0.3% pet	0.7 (1/139)	1.7 (1/59)	100	0 (0/80)	NA	0.424
Epoxy resin 1.0% pet	0.7 (1/137)	1.7 (1/59)	0	0 (0/78)	NA	0.431
Mercaptobenzothiazole 2% pet	0 (0/144)	0 (0/62)	NA	0 (0/82)	NA	NA
Thiuram mix 1% pet	0 (0/144)	0 (0/62)	NA	0 (0/82)	NA	NA
Colophonium 20% pet	0 (0/144)	0 (0/62)	NA	0 (0/82)	NA	NA
Neomycin sulfate 20% pet	0 (0/142)	0 (0/60)	NA	0 (0/82)	NA	NA
p-tert-Butylphenol formaldehyde resin 1% pet	0 (0/140)	0 (0/61)	NA	0 (0/79)	NA	NA
Budesonide 0.1% pet	0 (0/140)	0 (0/60)	NA	0 (0/80)	NA	NA
Mercapto mix 2% pet	0 (0/139)	0 (0/60)	NA	0 (0/79)	NA	NA
Tixocortol-21-pivalate 1% pet	0 (0/137)	0 (0/59)	NA	0 (0/78)	NA	NA
Disperse blue 106 1% pet	0 (0/137)	0 (0/59)	NA	0 (0/78)	NA	NA
Panthenol 5% pet	0 (0/131)	0 (0/57)	NA	0 (0/74)	NA	NA
Hydrocortisone-17-butyrate 0,1% pet	0 (0/110)	0 (0/43)	NA	0 (0/67)	NA	NA
Primin 0,01% pet	0 (0/97)	0 (0/38)	NA	0 (0/59)	NA	NA
Carba mix 3% pet	0 (0/95)	0 (0/45)	NA	0 (0/50)	NA	NA
Farnesol 5% pet	0 (0/91)	0 (0/36)	NA	0 (0/55)	NA	NA
Clioquinol 5% pet	0 (0/90)	0 (0/36)	NA	0 (0/54)	NA	NA
Textile dye mix 6.6% pet	0 (0/49)	0 (0/24)	NA	0 (0/25)	NA	NA
DMDM hydantoin 1% pet	0 (0/47)	0 (0/25)	NA	0 (0/22)	NA	NA
Lidocaine 15% pet	0 (0/45)	0 (0/22)	NA	0 (0/23)	NA	NA

Cosmetic series*						
Cocamidopropyl betaine 1% aq	2.3 (1/44)	0 (0/21)	NA	4.3 (1/23)	100	>0.99
Octyl gallate 0.25% pet	2.1 (1/47)	0 (0/22)	NA	4.0 (1/25)	100	>0.99
Phenoxyethanol 1% pet	2.0 (1/49)	0 (0/23)	NA	3.8 (1/26)	100	>0.99
Cetearyl alcohol 20% pet	1.9 (1/52)	0 (0/24)	NA	3.6 (1/28)	100	>0.99
Triethanolamine 2.5% pet	0 (0/56)	0 (0/25)	NA	0 (0/31)	NA	NA
Ethylenediaminetetraacetic acid 1% pet	0 (0/55)	0 (0/24)	NA	0 (0/31)	NA	NA
Chlorocresol 1% pet	0 (0/54)	0 (0/24)	NA	0 (0/30)	NA	NA
Oxybenzone 10% pet	0 (0/52)	0 (0/23)	NA	0 (0/29)	NA	NA
Sorbic acid 2% pet	0 (0/51)	0 (0/24)	NA	0 (0/27)	NA	NA
Chloroxyleneol 1% pet	0 (0/50)	0 (0/23)	NA	0 (0/27)	NA	NA
Hexahydro-1,3,5-tris-(2-hydroxyethyl)triazine (Grotan BK) 1% aq	0 (0/49)	0 (0/23)	NA	0 (0/26)	NA	NA
Chloroacetamide 0.2% pet	0 (0/49)	0 (0/23)	NA	0 (0/26)	NA	NA
Propylene glycol 5% pet	0 (0/48)	0 (0/23)	NA	0 (0/25)	NA	NA
Butyl hydroxy toluene 2% pet	0 (0/47)	0 (0/22)	NA	0 (0/25)	NA	NA
Triclosan 2% pet	0 (0/47)	0 (0/21)	NA	0 (0/26)	NA	NA
Abietic acid 10% pet	0 (0/46)	0 (0/22)	NA	0 (0/24)	NA	NA
Butylhidroxianisole 2% pet	0 (0/43)	0 (0/20)	NA	0 (0/23)	NA	NA

Numbers in parentheses represent the number of times that a positive reaction was found to allergen, divided by the total number of times that the allergen was tested. PRPP: Present Relevant per positives; P-value is result of differences between patients with and without atopic dermatitis groups. Statistically significant values were marked bold.

*Allergens patch tested <20 times were excluded to avoid non-inferential results (excluded ones are presented in supplementary information).

MCI/MI: methylchloroisothiazolinone/methylisothiazolinone; NA: not applicable.

Table 3. Top 20 most common sensitization prevalence rates for allergens from European baseline series and cosmetic series. Allergens are shown from most to least frequent.

Top 20 Total		Top 20 With atopic dermatitis		Top 20 Without atopic dermatitis	
1.	Nickel sulfate 5% pet	1.	Nickel sulfate 5% pet	1.	Nickel sulfate 5% pet
2.	MCI/MI 0.02% aq	2.	Hydroperoxide linalool 1/0,5% pet	2.	Benzisotiazolinona 0.1% pet
3.	Hydroperoxide linalool 1/0,5% pet	3.	Quaternium 15 1% pet	3.	Methylisothiazolinone 0.2% aq
4.	Methylisothiazolinone 0.2% aq	4.	MCI/MI 0.02% aq	4.	MCI/MI 0.02% aq
5.	Amerchol L-101 50% pet	5.	Parthenolide 0.1% pet	5.	Caine mix III 10% pet
6.	Cobalt chloride 1% pet	6.	Fragrance mix I 8% pet	6.	Hydroperoxide linalool 1/0,5% pet
7.	Benzisotiazolinona 0.1% pet	7.	Diazolidinyl urea 2% pet	7.	Sodium disulfite 1% pet
8.	Fragrance mix I 8% pet	8.	Amerchol L-101 50% pet	8.	p-Phenylenediamine 1% pet
9.	Caine mix III 10% pet	9.	Disperse Orange 3 1% pet	9.	Fragrance mix I 8% pet
10.	p-Phenylenediamine 1% pet	10.	2-Hydroxyethyl methacrylate 1% pet	10.	Amerchol L-101 50% pet
11.	Disperse Orange 3 1% pet	11.	Hidroperoxide of limonene 0.2/0.3% pet	11.	Cocamidopropyl betaine 1% aq*
12.	Formaldehyde 2% aq	12.	p-Phenylenediamine 1% pet	12.	Octyl gallate 0.25% pet*
13.	Parthenolide 0.1% pet	13.	Imidazolidinyl urea 2% pet	13.	Cobalt chloride 1% pet
14.	Diazolidinyl urea 2% pet	14.	Formaldehyde 2% aq	14.	Phenoxyethanol 1% pet*
15.	Quaternium 15 1% pet	15.	Compositae mix II 2.5% pet	15.	Formaldehyde 2% aq
16.	Sodium disulfite 1% pet	16.	1,2-Dibromo-2,4-dicyanobutane 0,3% pet	16.	Fragrance mix II 14% pet

17.	Cocamidopropyl betaine 1% aq*	17.	Caine mix II 10% pet	17.	Cetearyl alcohol 20% pet*
18.	N-Isopropyl-N'-phenyl-pphenylenediamine 0.1% pet	18.	N-Isopropyl-N'-phenyl-pphenylenediamine 0.1% pet	18.	Disperse Orange 3 1% pet
19.	Hidroperoxide of limonene 0.2/0.3% pet	19.	Sesquiterpene lactone mix 0.3% pet	19.	Lanolin alcohols 30% pet
20.	Lanolin alcohols 30% pet	20.	Epoxy resin 1.0% pet	20.	N-Isopropyl-N'-phenyl-pphenylenediamine 0.1% pet

Different top 20 allergens between with and without atopic dermatitis groups were marked bold.

*Allergens from Cosmetic series.

MCI/MI: methylchloroisothiazolinone/methylisothiazolinone.

Table 4. Top 20 most common relevant allergens from European baseline series and cosmetic series. Allergens are shown from most frequently relevant in number to least frequent.

Top 20 With atopic dermatitis		Top 20 Without atopic dermatitis	
1.	Quaternium 15 1% pet	1.	Benzisotiazolinona 0.1% pet
2.	MCI/MI 0.02% aq	2.	Hydroperoxide linalool 1/0,5% pet
3.	Parthenolide 0.1% pet	3.	p-Phenylenediamine 1% pet
4.	Fragrance mix I 8% pet	4.	Cocamidopropyl betaine 1% aq*
5.	Diazolidinyl urea 2% pet	5.	Octyl gallate 0.25% pet*
6.	2-Hydroxyethyl methacrylate 1% pet	6.	Phenoxyethanol 1% pet*
7.	Hidroperoxide of limonene 0.2/0.3% pet	7.	Cetearyl alcohol 20% pet*
8.	p-Phenylenediamine 1% pet	8.	Hydroxyisohexyl 3-cyclohexene carboxaldehyde (Iyral) 5% pet
9.	Imidazolidinyl urea 2% pet	9.	Propolis 10% pet
10.	Compositae mix II 2.5% pet	10.	Imidazolidinyl urea 2% pet
11.	1,2-Dibromo-2,4-dicyanobutane 0,3% pet	11.	Diazolidinyl urea 2% pet
12.	Caine mix II 10% pet	12.	Peru balsam 25% pet
13.	N-Isopropyl-N'-phenyl-pphenylenediamine 0.1% pet	13.	Methylisothiazolinone 0.2% aq
14.	Sesquiterpene lactone mix 0.3% pet	14.	MCI/MI 0.02% aq
15.	Methylisothiazolinone 0.2% aq	15.	Fragrance mix I 8% pet
16.	Potassium dichromate 0.5% pet	16.	Fragrance mix II 14% pet
17.	Parabens 12% pet	17.	Formaldehyde 2% aq
18.	Nickel sulfate 5% pet	18.	Nickel sulfate 5% pet
19.	Amerchol L-101 50% pet	19.	Lanolin alcohols 30% pet
20.	Hydroperoxide linalool 1/0,5% pet	20.	Cobalt chloride 1% pet

Different top 20 allergens between with and without atopic dermatitis groups were marked bold;

*Allergens from Cosmetic series.

MCI/MI: methylchloroisothiazolinone/methylisothiazolinone.

Figure 1. Purpose for the adapted baseline series for children with AD, with suspected ACD, difficult-to-control AD, new-onset dermatitis and prior to initiating systemic therapy for AD, or when no specific exposure is identified.

BASELINE SERIES FOR CHILDREN WITH ATOPIC DERMATITIS WITH RECOMMENDATION TO ACD EVALUATION		
Allergens	48 h	72-96h
1. Quaternium 15 1% pet		
2. MCI/MI 0.02% aq		
3. Parthenolide 0.1% pet		
4. Fragrance mix I 8% pet		
5. Diazolidinyl urea 2% pet		
6. 2-Hydroxyethyl methacrylate 1% pet		
7. Hydroperoxide of limonene 0.2/0.3% pet		
8. p-Phenylenediamine 1% pet		
9. Imidazolidinyl urea 2% pet		
10. Compositae mix II 2.5% pet		
11. 1,2-Dibromo-2,4-dicyanobutane 0.3% pet		
12. Caine mix II 10% pet		
13. N-Isopropyl-N'-phenyl-phenylenediamine 0.1% pet		
14. Sesquiterpene lactone mix 0.3% pet		
15. Methylisothiazolinone 0.2% aq		
16. Potassium dichromate 0.5% pet		
17. Parabens 12% pet		
18. Nickel sulfate 5% pet		
19. Amerchol L-101 50% pet		
20. Hydroperoxide linalool 1/0,5% pet		