SHORT REVIEW

Lessons from peculiar cases of anaphylaxis: why allergists should be prepared for the unexpected.

G. Liccardi ^{1*†}, M. Milanese ^{2*}, MB Bilò ^{3*}, MV Liccardi ^{4*}, D. Gargano ^{5†}, A. Giordano ^{6*}, F. Habetswallner ^{7*}, M. Lo Schiavo ^{8†}, F. Madonna ^{9†}, MC. Montera ^{8†}, G. Pace ^{10*}, A. Papa ^{11†}, A. Pedicini ^{12†} and P. Rogliani ^{1,10*}

On behalf of the multidisciplinary Working Groups “Stress-Asthma / Allergy”^{*} and “AAIITO Regione Campania”[†]

Data presented at AAIITO 2020 -On Line “Il Mese dell’Allergologia AAIITO” 9-26 October 2020.

¹ Postgraduate School of Respiratory Medicine. Department of Experimental Medicine, University of Rome "Tor Vergata", Rome, Italy

² Division of Pulmonology, S. Corona Hospital, Pietra Ligure, Italy.

³ Department of Clinical and Molecular Sciences, Marche Polytechnic University - Allergy Unit, Department of Internal Medicine, University Hospital Ospedali Riuniti - Ancona, Italy.

⁴ Psychologist

⁵ Allergy Unit. High Speciality “San Giuseppe Moscati” Hospital, Avellino

⁶ Postgraduate School of Internal Medicine. University of Salerno

⁷ Division of Neurophysiology “A. Cardarelli” Hospital, Naples, Italy.

⁸ Allergy and Clinical Immunology. “G. Fucito” Hospital and University Hospital, Salerno

⁹ Allergy Unit ASL (Sanitary District n°12), Caserta

¹⁰ Department of Experimental Medicine, Unit of Respiratory Medicine. University of Rome "Tor Vergata", Rome, Italy

¹¹ ASL (Sanitary District), Avellino

¹² Unit of Allergology. Division of Internal Medicine, "Fatebenefratelli" Hospital, Benevento

Word count:- 2572

Short title: Rare anaphylaxis.

Key words: adverse drug reaction, allergen exposure, anaphylaxis, bronchial asthma, Can f 5, COVID-19 vaccines, dog allergen, hypersensitivity, intimate behaviour, oxytocine, polyethylene glycol (PEG), psychological stress,

Summary statement: sometimes anaphylaxis can occur with uncommon or peculiar characteristics which could delay diagnosis and therapeutic treatment. We aimed to describe less accounted / difficult-to-approach shapes of anaphylaxis to facilitate clinicians to suspect these severe reactions even in uncommon conditions.

Authorship: All authors contributed equally in the writing and revision of the manuscript.

Conflict of interest and financial resources. All authors declare that they have no conflict of interest and that the study has been carried out without any financial support.

Corresponding author

Gennaro Liccardi, MD

-Postgraduate School of Respiratory Medicine. Department of Experimental Medicine, University of Rome "Tor Vergata", Rome, Italy

Phone : +39 081 5780554; E-mail: gennaro.liccardi51@gmail.com

Summary

Anaphylaxis is the most severe systemic hypersensitivity reaction, it can be caused by a number of well identified triggers such as foods, drugs, stinging insects and facilitated by predisposing clinical conditions. However, sometimes anaphylaxis shows up with uncommon or peculiar characteristics which could delay diagnosis and therapeutic treatment. In this report we aimed to describe less accounted / difficult-to-approach shapes of anaphylaxis to facilitate clinicians to suspect these severe reactions even in uncommon conditions. We choose to present data on anaphylaxis regarding: simulation, mode of exposure to sensitizing agents, pregnancy, exposure to animals, intimate behaviour, psychological stress and other situations.

Introduction

Anaphylaxis is the most severe systemic hypersensitivity reaction, it involves multiple organ systems and can be deadly (1,2). Anaphylaxis can be caused by a number of well identified triggers such as foods, drugs, stinging insects and clinical conditions (1,2). However, sometimes anaphylaxis shows up with uncommon or peculiar characteristics which could delay diagnosis and therapeutic treatment. In this brief report we aimed to introduce less accounted / difficult-to-approach shapes of anaphylaxis to facilitate clinicians to suspect these severe reactions even in uncommon conditions.

~~Atypical presentation of~~ Simulated anaphylaxis

In rare occasions, it has been reported that anaphylaxis can be mimicked by other clinic conditions (3) such as a diverticular perforation associated with colo-urachal fistula (4) and a massive subcutaneous emphysema presenting as anaphylaxis (5). We have recently shown that in obstetric surgery oxytocin/vasopressin can induce some side effects (negative inotropic and chronotropic effects, low blood pressure, vasodilatation) which can simulate a cardiac anaphylaxis and consequent intraoperative diagnostic problems and risks for the patient (6).

Anaphylaxis and modality mode of exposure

Inhalation-mode

Inhalation of the offending allergens is an uncommon cause of anaphylaxis. Inhalant allergens such as those of some animals (guinea pig, horse, deer and rabbit) (7,8) and grass pollens (9) can induce anaphylaxis. Food allergens, particularly from milk, in aerosolized form has been associated to anaphylaxis in a dairy worker (10) or in children as component of a medication aerosol (11). Anaphylaxis after occupational inhalation of Cefuroxime in a nurse has been also described (12).

Contact-mode

It has been published only five cases of anaphylaxis induced by skin contact with milk or milk-derived foods; some indications on the risk factors to identify susceptible individuals are shown in **table 1**(13). Recently a fatal reaction has been registered in a child at school;

(14) and another recent case report showed a generalized reaction induced by a cutaneous exposure to milk proteins in a colostrum-based cream (15).

Skin prick test (SPT)-related anaphylactic reactions can also be ascribed to the contact mode of exposure, those induced by the use of foods or drugs being the most frequent in clinical practice (16). On the contrary-anaphylaxis produced by SPT with standard aero-allergens is extremely uncommon. We described a rare case of systemic life-threatening reaction after a routine SPT in a patient with high allergic sensitization to parietaria and grasses (17). In the same article we reported a multicentric evaluation of the frequency of SPT-related-anaphylaxis in Italy: our case was the only one among 55.105 patients and 624.500 allergens tested (17). Possible risk factors for SPT-related anaphylaxis are similar to those described in table 1. Moreover, in children we suggest to avoid the contemporary use of inhalant and food allergens, and the use of intradermal skin testing, to minimize the number of allergens to be tested during the SPT procedure, and to check the subjects for at least 20 min after the end of SPT. In some cases, it should be useful to consider the possibility of diluting allergenic extracts before their use (18,19). It is important to underline that the need to carry out "in vivo" tests (e.g. SPTs, prick by prick) for the diagnosis of food allergy in highly sensitized individuals has been reduced by the growing use and utility of modern laboratory procedures such as Component Resolved Diagnostics (20, 21).

Anaphylaxis and pregnancy

Anaphylaxis (A) is considered uncommon during pregnancy. However, among potential risk factors such as age and history of allergy, cesarean delivery could be recognized as the main culprit condition because the necessity of using anaesthesia, surgical procedures and drugs (22). An amoxicillin-induced anaphylaxis at the end of pregnancy determined newborn death by hypoxic-ischaemic cerebral injuries (23).

As previously reported, delivery, especially when carried out by cesarean section, can determine dangerous systemic reactions. Oxytocin constitute a complex cause of systemic / bronchial reactions as well as equivocal side effects-related- symptoms (negative inotropic and chronotropic effects, low blood pressure, vasodilatation) that mimic cardiac (A) anaphylaxis (6, 24). Another insidious effect derived by a demonstrated homology in the protein sequence of oxytocin and latex allergens Hev b 7.01 and Hev b 7.02, (25).

We have previously described two life-threatening anaphylactic reactions with onset a few minutes after the infusion of oxytocin in two women sensitized to latex allergens during caesarian section under spinal anaesthesia in the delivery room (26). Moreover, it has been also demonstrated that oxytocin, under pro-inflammatory cytokines stimulation, may induce contraction of smooth muscle and airway narrowing suggesting that oxytocin serves as a bronchoconstrictor (27). These data suggest that inflammatory conditions of airways such as those found in asthmatic women (especially those with severe asthma) might constitute an independent (from A anaphylaxis) risk factor for airway obstruction after infusion of oxytocin during delivery. The role of oxytocin receptors could also explain the well-known worsening of asthma control in about one-third of pregnant women suffering from asthma. These findings suggest a particular attention in managing delivering women suffering from latex allergy and bronchial asthma. An accurate anamnestic clinical and diagnostic evaluation, a latex-free setting, the use of oxytocin-alternative agents and in case of asthma, a drug premedication are likely to reduce the risk of anaphylactic or airway-obstructive reactions in these women (24) (**Figure 1**).

Another extremely uncommon condition for anaphylaxis is lactation (28). It is not clear the pathogenesis, a possible role of hormones has been suggested (the withdrawal of the stabilising effect of progesterone on mast cells) but also the influence of others (prolactin, oxytocin, adrenocorticotrophic hormone and corticotropin-releasing hormone) (29).

Finally, some case reports have shown that bovine serum albumin may be a causative agent in severe (A) anaphylaxis after standard intrauterine insemination or in vitro fertilization if added to the culture medium of spermatozooids (30)

Anaphylaxis and animal exposure

Although respiratory allergy is a common consequence of animal allergens inhalation, in rare occasions a clear clinical picture of anaphylaxis developed after inhalation of guinea pig, horse, rabbit and deer epithelia (7,8).

Another possible modality of inducing anaphylaxis after exposure to animals is their bite. In some rare cases the bites were inflicted by domestic or wild animals such as horse, gerbil, mouse (31-33).

However, the risk of anaphylaxis from animal bites is relatively more common in individuals working with laboratory animals, particularly rodents (34).

Another topic related to animals regards some food-derived animal substances that can be used in medications at various stages of the manufacturing process. Caglayan-Sozmen et al. (35) review the possible roles of medications which may contain egg, red meat, gelatin, and fish allergens in allergic

reactions in children with food allergy. The risk of reactions to hidden milk allergens in drugs is particularly important. In fact, dry-powder inhalers may contain lactose as excipient, this product can be rarely contaminated with milk proteins and, consequently, it may induce allergic reactions in patients with cow's milk allergy (36, 37). Reports have described immediate hypersensitivity reactions to methylprednisolone sodium succinate 40 mg injection, a formulation that contains lactose as excipient (38). Some cases of anaphylaxis after receiving diphtheria-tetanus-pertussis vaccine injection (39) or some probiotics (40) in children allergic to milk have been reported. Among food-derived animal substances also gelatin can induce systemic reactions being contained in medications like plasma volume expanders, erythropoietin, hemostatic products (41), as well as in some vaccines (42,43). Although dogs can induce respiratory allergy, we would like to cite the potential role of service dogs in certain clinical conditions at high risk of anaphylaxis such as surgical interventions in subjects with high allergic risk. Tew & Taicher documented the first report of a service dog used to detect mast cell mediator release in patients with mastocytosis (44). This service dog was used not only in a family-centered care model, but also as an additional perioperative monitor to predict the eventual occurrence of perioperative mediator release (44).

Anaphylaxis and intimate behaviour

Although rare, the contact with allergens can also be due to a direct contact between two individuals, one of whom carries on his/her body the allergen potentially dangerous for the other individual (or can carry traces of food or drugs in biological fluids) (Figure 2). This can easily happen during intimate behaviours, such as kisses or sexual intercourses. This topic may be underestimated and under-reported, due to its delicate nature, involving a very intimate aspect of the life (45).

The mechanism by which kissing may induce local or, in some cases, generalized reactions in sensitized individuals is the passive transport of allergenic molecules through saliva, skin or oral mucosa, and the consequent contact of these allergens with the skin or mucosae of the sensitized subjects. The severity of allergic symptoms likely depends on the type of kissing, as the contact with the 'unprotected' oral mucosa may account for a greater penetration of allergens in comparison to intact skin (45).

Foods are the most frequent cause of allergic reactions following a kiss from an individual who have eaten an allergenic food to a patient sensitized to the same allergen (46). Peanuts, walnuts and other tree nuts are the most common foods involved, followed by fruits or vegetables such as apple, carrot, kiwi, fish, shellfish, and milk (47,48). There are only few reports of oral allergy syndrome and consequent generalized urticaria in drug-sensitized individuals, a few minutes after passionate kisses given by their partners who had previously used the drug particularly beta lactams (49,50). Foods (such as Brazil nut) can also penetrate into seminal fluid and induce-after vaginal intercourse (51); seminal fluid can also carry antibiotics (such as amoxicillin) and trigger-anaphylaxis in an amoxicillin-sensitized woman after oral sex (52).-Anaphylaxis after anal intercourse but tolerance after vaginal intercourse has been also described (53).

Seminal plasma hypersensitivity (SPH) is characterized by immediate local or systemic postcoital symptoms following mucosal exposure to seminal fluid. Systemic SPH is a type I, immediate, IgE-mediated hypersensitivity reaction with the well-known symptoms (54).

The prostate-specific antigen (PSA), a 33–34 kDa glycoprotein, better known for its use in monitoring prostate cancer in men, was in fact identified as a major allergen in human seminal fluids. It was found that PSA carries high homology to the canine prostatic kallikrein, which was identified as Can f 5. The clinical relevance of this cross-reactivity was confirmed in a few cases, and investigators have hypothesized that patients who experienced SPH after their first unprotected intercourse could have been sensitized by previous exposure to dogs (54).

We have recently shown the increasing relevance of allergic sensitization to Can f 5 in North East Italy considering both the number of sensitized individuals (69.02% among 268 dog-sensitized ones, and 57.92% as exclusive sensitization) and the level of sensitization (77.6% of individuals showed medium or high values of specific IgE according to manufacturer's specifications) (55). Moreover, our Italian multicenter study has shown that a prevalent exposure to a male dog represents a risk factor for the presence of Can f 5 monosensitization (56). The results of this study emphasize the need of an adequate diagnosis and management of patients suffering from dog allergy, especially those with relevant clinical symptoms following dog exposure in order to prevent systemic EAH (57).

Anaphylaxis and psychological stress

The relationship between anaphylaxis and psychological stress (PS) is an intriguing puzzle, the question being: can the former be the cause or only consequence of the latter or both? It is well documented that (PS) may enhance allergic diseases such as asthma, allergic rhinitis, and atopic dermatitis through several complex mechanisms like mast cell activation and mediator release, inflammation and imbalance of the autonomic system (58-60). Stress can activate the hypothalamic-pituitary-adrenal axis through the release of corticotropin-releasing hormone (CRH). Theoharides et al. have shown that CRH secreted under stress stimulates mast cell degranulation through activation of CRH receptor-1 (CRHR-1) (61). This effect is augmented by other neuropeptides also released by stress, such as substance P (SP) and neurotensin (NT). In fact, both NT and SP induce the expression of functional CRHR-1 (61). Moreover, the same authors have shown that CRH induces the expression of high-affinity IgE receptor (FcεRI) and augments allergic stimulation of human mast cells (62). To the best of our knowledge, only two case reports have clearly suggested a relationship between a PS situation and the onset of anaphylaxis (63,64). The diagnosis of such anaphylaxis has been made essentially through anamnestic criteria and exclusion of other most common triggering agents / situations. We believe that PS alone or as a relevant co-factor could induce anaphylaxis more frequently than expected if we consider the aforementioned mechanisms and the high presence of PS in the general population. On the contrary, the role/onset of PS in patients surviving anaphylaxis episodes has been better explored. Baiardini et al. (65) studied quality of life and well-being in patients with drug-induced anaphylaxis. Two validated tools were used: The Drug Hypersensitivity Quality-of-Life Questionnaire (DrHy-Q) and the Psychological General Well-Being Index (PGWBI). Compared with the Italian reference population, patients had a significantly reduced PGWBI total and domain score. The authors highlighted for the first time impaired Health Related Quality of Life (HRQoL) and distress commonly feature in survivors to anaphylactic reactions to drug. Recently Lee et al. (66) have evaluated Posttraumatic Stress Disorder (PTSD) arising after-anaphylaxis in adults, by investigating the psychosocial burden of recent-anaphylaxis in Korean adults. The results of this study indicated that patients who experienced anaphylaxis were likely to develop psychiatric disorders, such as PTSD, anxiety, and depression. The development of PTSD did not appear to depend on the severity of anaphylaxis, although patients with higher Impact of Event Scale-Revised-Korean version (IES-R-K), scores had more severe anxiety and depression. These and previous considerations suggest that the general management (diagnosis / treatment) of anaphylaxis should include a psychological / psychiatric evaluation.

Others

Anaphylaxis (A) after the use of polyethylene glycol (PEG), a water-soluble, organic compound included in a wide variety of products has been described. In the healthcare setting, it is a common ingredient in medications and procedural agents (eg. for performing ultrasound scans) (67). Anaphylaxis (A) to chlorhexidine (a synthetic bisbiguanide antiseptic agent) has been reported particularly in the perioperative and medical procedural settings (68). Rare cases of (A) anaphylaxis have been reported also with the use of anti-anaphylaxis agents such as cetirizine and chlorpheniramine (69,70). Another controversial topic is the possible induction of anaphylaxis by vaccines used to prevent contagious infections, many patients refusing vaccinations for this fear. Mc Neil et al. (71) reported only 33 vaccine-triggered anaphylaxis cases that occurred after 25,173,965 vaccine doses (against different infectious agents), the rate of anaphylaxis being 1.31 (95% CI, 0.90-1.84) per million vaccine doses. Interest in this issue has increased significantly with the recent start of mass vaccination against the Sars-CoV-2 virus. Preliminary reports have been published on anaphylactic events occurring after the administration of the first Sars-CoV-2 available vaccines. Twenty-one cases were determined to be anaphylaxis (a rate of 11.1 per million doses administered) after the use of Pfizer-BioNTech COVID-19 Vaccine (72), while 10 cases were determined to be anaphylaxis (a rate of 2.5 anaphylaxis cases per million Moderna COVID-19 vaccine doses administered) (73). The aforementioned polyethylene glycol (PEG) has been indicated as potential agent of anaphylaxis in these vaccines being an excipient (74,75,76).

Conclusions

Data reported from our previous studies and from a review of the Literature show that sometimes anaphylaxis can occur with uncommon or peculiar characteristics which could delay diagnosis and therapeutic treatment. We hope that the description of less accounted / difficult-to-approach shapes of anaphylaxis could facilitate clinicians to suspect these severe reactions even in less common clinic presentations.

Acknowledgements

We thank the biologist Dr. Paola Derra and miss Lucia Franzese for technical assistance in the preparation of this manuscript.

References

1. Bilò MB, Martini M, Tontini C, Corsi A, Antonicelli L. Anaphylaxis. Eur Ann Allergy Clin Immunol. 2021; 53: 4-17
2. Muraro A, Roberts G, Worm M, Bilò MB, Brockow K, Fernández Rivas M, Santos AF, Zolkipli ZQ, Bellón A, Beyer K, Bindslev-Jensen C, Cardona V, Clark AT, Demoly P, Dubois AE, DunnGalvin A, Eigenmann P, Halken S, Harada L, Lack G, Jutel M, Niggemann B, Ruëff F, Timmermans F, Vlieg-Boerstra BJ, Werfel T, Dhimi S, Panesar S, Akdis CA, Sheikh A; EAACI Food Allergy and

- Anaphylaxis Guidelines Group. Anaphylaxis: guidelines from the European Academy of Allergy and Clinical Immunology. *Allergy*. 2014; 69:1026-45.
- 3.Yuson C, Kette F, Hissaria P, Smith W. Rare mimic of recurrent anaphylaxis. *BMJ Case Rep* 2019; 12: e228173.
- 4.Yang D, Pearson D, Smith D. A rare case of a diverticular perforation associated with colo-urachal fistula presenting as anaphylaxis. *Ann R Coll Surg Engl*. 2020;102: e51-e53.
5. Earley A, Watkins T, Forde N. Massive subcutaneous emphysema mimicking anaphylaxis - pathological and radiological correlations. *Forensic Sci Med Pathol*. 2019; 15: 603-606.
6. Liccardi G, Calzetta L, Salzillo A, Puxeddu E, Rogliani P. Relationship between oxytocin/vasopressin and latex in obstetric surgery: how to recognize (and prevent) allergic reactions and differentiate them from side effects? *J Allergy Clin Immunol Pract*. 2017; 5: 873.
7. Cavkaytar Ö, Soyer Ö, Şekerel BE. A rare cause of aeroallergen-induced anaphylaxis: horse allergy. *Turk J Pediatr*. 2014; 56: 437-39.
8. Tang R, Li H. Anaphylaxis induced by inhaling guinea pig allergen. *Chin Med J (Engl)*. 2020 Aug 24. doi: 10.1097/CM9.0000000000000971.
9. Haluk Akar H, Akif Goktas M, Tahan F. Grass pollen triggered anaphylaxis in an adolescent boy. *Eur Ann Allergy Clin Immunol*. 2015; 47: 20-21.
10. Quirantes Sierra B, Lara Jiménez A, Skodova M. Sensitization to cow's milk protein in a dairy worker. *Occup Med (Lond)*. 2017; 67: 579-580.
11. Santoro A, Andreozzi L, Ricci G, Mastrorilli C, Caffarelli C. Allergic reactions to cow's milk proteins in medications in childhood. *Acta Biomed*. 2019; 90(3-S): 91-93
12. Merget R, Sander I, Fartasch M, van Kampen V, Röseler S, Merk H, Wurpts G, Raulf M, Brünig Occupational generalized urticaria and anaphylaxis after inhalation of cefuroxime in a nurse. *Am J Ind Med*. 2018; 61: 261-266.
13. Liccardi G, De Falco F, Gilder JA, D'Amato M, D'Amato G. Severe systemic allergic reaction induced by accidental skin contact with cow milk in a 16-year-old boy. A case- report. *J Investig Allergol Clin Immunol*. 2004; 14: 168-71.
14. Foong RX, Turner PJ, Fox AT. Fatal anaphylaxis due to transcutaneous allergen exposure: An exceptional case. *J Allergy Clin Immunol Pract*. 2020; 8: 332-333
15. Porcaro F, Caminiti L, Crisafulli G, Guglielmo F, Giovanni Battista Pajno GB. Anaphylaxis to cutaneous exposure to bovine colostrum-based cream. *Asian Pac J Allergy Immunol* 2019; 37: 9-11
16. Liccardi G, D'Amato G, Canonica GW, Salzillo A, Piccolo A, Passalacqua G. Systemic reactions from skin testing: literature review. *J Investig Allergol Clin Immunol*. 2006;16: 75-8.

- 373 17. Liccardi G, Salzillo A, Spadaro G, Senna G, Canonica WG, D'amato G, Passalacqua G.
374 Anaphylaxis caused by skin prick testing with aeroallergens: Case report and evaluation of the risk in
375 Italian allergy services. *J Allergy Clin Immunol*. 2003; 111:1410-2.
376
- 377 18. Liccardi G, Salzillo A, D'Amato G. The risk of generalized allergic reactions to skin prick testing
378 may be higher in poly-sensitized individuals and in those who underwent SPT with fresh foods. *Pediatr*
379 *Allergy Immunol*. 2010; 21: 656-57
380
- 381 19. Liccardi G, Calzetta L, Milanese M, Apicella G, Rogliani P. The possible concomitant use of
382 aeroallergen and food panels for skin prick testing might enhance the risk of generalized allergic
383 reactions in children. *Turk J Pediatr*. 2019; 61: 815-816.
384
- 385 20. Matricardi PM, Kleine-Tebbe J, Hoffmann HJ, Valenta R, Hilger C, Hölzle S, Agache I, Asero
386 R, Ballmer-Weber B, Barber D, Beyer K, Biedermann T, Bilò MB, Blank S, Bohle B, Bosshard PP,
387 Breiteneder H, Brough HA, Caraballo L, Caubet JC, Cramer R, Davies JM, Douladiris N, Ebisawa M,
388 Elgenmann PA, Fernandez-Rivas M, Ferreira F, Gadermaier G, Gatz M, Hamilton RG, Hawranek T,
389 Hellings P, Hoffmann-Sommergruber K, Jakob T, Jappe U, Jutel M, Kamath SD, Knol EF, Korosec P,
390 Kuehn A, Lack G, Lopata AL, Mäkelä M, Morisset M, Niederberger V, Nowak-Węgrzyn AH,
391 Papadopoulos NG, Pastorello EA, Pauli G, Platts-Mills T, Poon D, Poulsen LK, Raulf M, Sastre J,
392 Scala E, Schmid JM, Schmid-Grendelmeier P, van Hage M, van Ree R, Vieths S, Weber R, Wickman
393
- 394 21. M, Muraro A, Ollert M. et al. EAACI Molecular Allergy User's Guide. *Pediatr Allergy*
395 *Immunol*. 2016;27 Suppl 23:1–250.
396
- 397 22. McCall SJ, Kurinczuk JJ, Knight M. Anaphylaxis in pregnancy in the United States: risk factors
398 and temporal trends using national routinely collected data. *J Allergy Clin Immunol Pract*. 2019; 7:
399 2606-2612
- 400 23. Berenguer A, Couto A, Brites V, Fernandes R. Anaphylaxis in pregnancy: a rare cause of neonatal
401 mortality. *BMJ Case Rep*. 2013; 2013:bc2012007055
- 402 24. Liccardi G, Bilò M, Mauro C, Salzillo A, Piccolo A, D'Amato M, Liccardi A, D'Amato G.
403 Oxytocin: an unexpected risk for cardiologic and broncho-obstructive effects, and allergic reactions in
404 susceptible delivering women. *Multidiscip Respir Med*. 2013; 8: 67.
- 405 25. Ogata J, Minami K: Synthetic oxytocin and latex allergy. *Br J Anaesth* 2007; 98: 845–846.
- 406 26. Liccardi G, Bilò MB, Mauro C, Salzillo A, Piccolo A, D'Amato M, D'Amato G. Oxytocin: a likely
407 underestimated risk for anaphylactic reactions in delivering women sensitized to latex. *Ann Allergy*
408 *Asthma Immunol*. 2013; 110: 465-6.
- 409 27. Amrani Y, Svea F, Huang C, Li K, Liu V, Jain D, Keslacy S, Sims MW, Baidouri H, Cooper PR,
410 Zhao H, Siddiqui S, Brightling CE, Griswold D, Li L, Panettieri RS Jr: Expression and activation of the
411 oxytocin receptor in airway smooth muscle cells: Regulation by TNFalpha and IL-13. *Respir Res* 2010;
412 11: 104.
413
- 414 28. Villalta D, Martelli P. A case of breastfeeding anaphylaxis. *Eur Ann Allergy Clin Immunol* 2007;
415 39: 26-27

29. Shawkat E, Hussain N, Myers JE, Gillham J, Helbert M. Breast milk: friend or foe? *BMJ Case Rep.* 2011; 2011:bcr0120113766
30. Pagán JA, Postigo I, Rodríguez-Pacheco JR, Peña M, Guisantes JA, Martínez J. Bovine serum albumin contained in culture medium used in artificial insemination is an important anaphylaxis risk factor. *Fertil Steril.* 2008; 90: 2013.e17-9.
31. Guida G, Nebiolo F, Heffler E, Bergia R, Rolla G. Anaphylaxis after a horse bite. *Allergy.* 2005; 60: 1088-89
32. Watson J, Schobitz E, Davis J. Gerbil bite anaphylaxis. *Am J Emerg Med.* 2018; 36: 171.e5-171.e6.
33. Thewes M, Rakoski J, Ring J. Anaphylactic reaction after a mouse bite in a 9-year-old girl. *Br J Dermatol.* 1999; 141: 179
34. Stave GM, Lee EH, Darcey DJ. Laboratory Animal Bite Anaphylaxis: A National Survey: Part 1: Case Series and Review of the Literature. *J Occup Environ Med.* 2017; 59: 728-738.
35. Caglayan-Sozmen S, Santoro A, Cipriani F, Mastrorilli C, Ricci G, Caffarelli C. Hazardous Medications in Children with Egg, Red Meat, Gelatin, Fish, and Cow's Milk Allergy. *Medicina (Kaunas).* 2019; 55: 501
36. Robles J, Motheral L. Hypersensitivity reaction after inhalation of a lactose-containing dry powder inhaler. *J Pediatr Pharmacol Ther.* 2014; 19: 200-11.
37. Nowak-Węgrzyn A, Shapiro GG, Beyer K, Bardina L, Sampson HA. Contamination of dry powder inhalers for asthma with milk proteins containing lactose. *J Allergy Clin Immunol.* 2004; 113: 558-60.
38. Porcaro F, Paglietti MG, Diamanti A, Petreschi F, Schiavino A, Negro V, Pecora V, Fiocchi A, Cutrera R. Anaphylactic shock with methylprednisolone sodium succinate in a child with short bowel syndrome and cow's milk allergy. *Ital J Pediatr.* 2017; 43:104.
39. Kattan JD, Konstantinou GN, Cox AL, Nowak-Węgrzyn A, Gimenez G, Sampson HA, Sicherer SH. Anaphylaxis to diphtheria, tetanus, and pertussis vaccines among children with cow's milk allergy. *J Allergy Clin Immunol.* 2011;128: 215-18.
40. Martín-Muñoz M, Fortuni M, Caminoa M, Belver T, Quirce S, Caballero T. Anaphylactic reaction to probiotics. Cow's milk and hen's egg allergens in probiotic compounds. *Pediatr Allergy Immunol.* 2012; 23: 778-84.
41. Lied GA, Lund KB, Storaas T. Intraoperative anaphylaxis to gelatin-based hemostatic agents: a case report. *J Asthma Allergy.* 2019; 12: 163-167
42. Miller CK, Mendoza JC, Coop CA. Anaphylaxis to MMR Vaccine Mediated by IgE Sensitivity to Gelatin. *Mil Med.* 2020; 185: e1869-e1871

- 458 43. Lin SF, Jiang PL, Tsai JS, Huang YY, Lin SY, Lin JH, Liu DZ. Surface assembly of poly(I:C) on
459 polyethyleneimine-modified gelatin nanoparticles as immunostimulatory carriers for mucosal antigen
460 delivery. *J Biomed Mater Res B Appl Biomater*. 2019; 107: 1228-1237.
461
- 462 44. Tew S, Taicher BM. A dog is a doctor's best friend: The use of a service dog as a pet operative
463 assistant. *Case Rep Pediatr*. 2016; 2016: 9013520
- 464 45. Liccardi G, Caminati M, Senna G, Calzetta L, Rogliani P. Anaphylaxis and intimate behaviour.
465 *Curr Opin Allergy Clin Immunol*. 2017; 17: 350-355
- 466 46. Eriksson NE, Moller C, Werner S, et al. The hazards of kissing when you are food allergic. *J Invest*
467 *Allergol Clin Immunol* 2003; 13:149–154.
468
- 469 47. Wuthrich B. Oral allergy syndrome to apple after a lover's kiss. *Allergy* 1997; 52:236.
470
- 471 48. Mancuso G, Berdondini RM. Oral allergy syndrome from kiwi fruit after a lover's kiss. *Contact*
472 *Dermatitis* 2001; 45:41.
473
- 474 49. Liccardi G, D'Amato G. Drugs may also induce allergic reactions in sensitized individuals through
475 passionate kissing. *J Allergy Clin Immunol* 2007; 119:755
476
- 477 50. Liccardi G, Gilder J, D'Amato M, D'Amato G. Drug allergy transmitted by passionate kissing.
478 *Lancet*. 2002; 359: 1700.
- 479 51. Bansal AS, Chee R, Nagendran V, Warner A, Hayman G. Dangerous liaison: sexually transmitted
480 allergic reaction to Brazil nuts. *J Investig Allergol Clin Immunol*. 2007; 17:189-91
481
- 482 52. Gómez Caballero N, Almenara S, Tévar Toro A, Horga de la Parte JF. Anaphylaxis probably
483 induced by transfer of amoxicillin via oral sex. *BMJ Case Rep*. 2019;12: e227398.
484
- 485 53. Martí-Garrido J, López-Salgueiro R, Bartolomé-Zavala B, Perales-Chordá C, de Rojas DH.
486 Anaphylaxis after anal intercourse with tolerance by vaginal route. *Ann Allergy Asthma Immunol*.
487 2019;122: 346-347
488
- 489 54. Ghosh D, Bernstein JA. Systemic and localized seminal plasma hypersensitivity patients exhibit
490 divergent immunologic characteristics. *J Allergy Clin Immunol* 2014;134: 969–972.
- 491 55. Villalta D, Milanese M, Da Re M, Sabatino G, Sforza M, Calzetta L, Liccardi G. Frequency of
492 allergic sensitization to Can f 5 in North East Italy. An analysis of 1403 ISACs 112 (Component
493 Resolved Diagnosis) collected retrospectively. *Eur Ann Allergy Clin Immunol*. 2019; 51:186-189.
- 494 56. Liccardi G, Calzetta L, Bilò MB, Brusca I, Cecchi L, Costantino MT, Da Re M, Buzzulini F, Giani
495 M, Lombardi C, Meneguzzi G, Meriggi A, Milanese M, Onida R, Patella V, Pignatti P, Quercia O,
496 Ridolo E, Sabatino G, Sacerdoti C, Savi E, Scala E, Steinhilber G, Villalta D, Passalacqua G, Rogliani
497 P. A prevalent exposure to male dog is a risk factor for exclusive allergic sensitization to Can f 5: An
498 Italian multicenter study. *J Allergy Clin Immunol Pract*. 2020; 8: 2399-2401

499 57. Liccardi G, Calzetta L, Salzillo A, Apicella G, Piccolo A, Di Maro E, Rogliani P. Dog allergy: can
500 a prevalent or exclusive sensitization to Can f 5 be considered a lucky or negative event in real life?
501 Eur Ann Allergy Clin Immunol. 2018;50: 283-285
502

503 58. Heffner KL, Kiecolt-Glaser JK, Glaser R, et al., Stress and anxiety effects on positive skin test
504 responses in young adults with allergic rhinitis, Ann. Allergy Asthma Immunol. 2014; 113: 13–18.
505

506 59. Lu Y, Ho R, Lim TK, et al., Neuropeptide Y may mediate psychological stress and enhance TH2
507 inflammatory response in asthma, J. Allergy Clin. Immunol. 2015;135; 1061–1067
508

509 60. Gupta K, Harvima IT Mast cell-neural interactions contribute to pain and itch. Immunol Rev. 2018
510 Mar;282(1):168-187
511

512 61. Theoharides TC, Singh LK, Boucher W, Pang X, Letourneau R, Webster E, Chrousos G.
513 Corticotropin-releasing hormone induces skin mast cell degranulation and increased vascular
514 permeability, a possible explanation for its pro-inflammatory effects. Endocrinology. 1998; 139: 403-
515 413.
516

517 62. Asadi S, Theoharides TC. Corticotropin-releasing hormone and extracellular mitochondria augment
518 IgE-stimulated human mast-cell vascular endothelial growth factor release, which is inhibited by
519 luteolin. J Neuroinflammation.2012; 9:85.
520

521 63. Alevizos M, Karagkouni A, Kontou-Fili K, Theoharides TC. A probable case report of stress-
522 induced anaphylaxis. Ann Allergy Asthma Immunol. 2014;112: 383-84
523

524 64. Larenas-Linnemann DE, Costa-Domínguez ML, Creticos PS. Acute emotional stress proposed
525 as a risk factor for anaphylaxis in patients receiving allergen immunotherapy. Ann Allergy Asthma
526 Immunol. 2020;124: 314-317

527 65. Baiardini I, Gaeta F, Molinengo G, Laido F, Canonica GW, Romano A. Quality-of-life issues in
528 survivors to anaphylactic reactions to drugs. Allergy. 2015; 70: 877-79

529 66. Lee Y, Chang HY, Kim SH, Yang MS, Koh YI, Kang HR, Choi JH, Kim CW, Park HK, Kim JH,
530 Nam YH, Kim TB, Hur GY, Jung JW, Park KH, Kim MA, Kim J, Yoon J, Ye YM. A Prospective
531 Observation of Psychological Distress in Patients With Anaphylaxis. Allergy Asthma Immunol Res.
532 2020;12: 496-506

533 67. Jakubovic BD, Saperia C, Sussman GL. Anaphylaxis following a transvaginal ultrasound . Allergy
534 Asthma Clin Immunol. 2016; 12: 3

535 68. Chiewchalewong C, Sompornrattanaphan M, Wongsas C, Thongngarm T. Chlorhexidine Allergy:
536 Current Challenges and Future Prospects , J Asthma Allergy. 2020; 13: 127-133

537 69. Afonso N, Shetgaonkar P, Dang A, Rataboli PV. Cetirizine-induced anaphylaxis: a rare adverse
538 drug reaction. Br J Clin Pharmacol. 2009; 67: 577-8.

70. Lee SH, Lee Y, Woo SD, Doo KE, Ha CY, Lee YH, Ye YM. Chlorpheniramine-induced anaphylaxis: Two case reports and a retrospective review of pharmacovigilance database. *Medicine (Baltimore)*. 2019; 98: e18369.
71. McNeil MM, Weintraub ES, Duffy J, Sukumaran L, Jacobsen SJ, Klein NP, Hambidge SJ, Lee GM, Jackson LA, Irving SA, King JP, Kharbanda EO, Bednarczyk RA, DeStefano F. Risk of anaphylaxis after vaccination in children and adults. *J Allergy Clin Immunol*. 2016; 137: 868-78.
72. Allergic Reactions Including **Anaphylaxis** After Receipt of the First Dose of Pfizer-BioNTech **COVID-19 Vaccine** - United States, December 14-23, 2020. CDC COVID-19 Response Team; Food and Drug Administration. *MMWR Morb Mortal Wkly Rep*. 2021;70: 46-51.
73. Allergic Reactions Including Anaphylaxis After Receipt of the First Dose of Moderna COVID-19 Vaccine - United States, December 21, 2020-January 10, 2021. CDC COVID-19 Response Team; Food and Drug Administration. *MMWR Morb Mortal Wkly Rep*. 2021; 70:125-129.
74. Garvey LH, Nasser S. Br J Anaphylaxis to the first COVID-19 vaccine: is polyethylene glycol (PEG) the culprit? *Anaesth*. 2020 Dec 17: S0007-0912(20)3009-6. doi: 10.1016/j.bja.2020.12.020.
75. Caballero ML, Quirce S. Excipients as potential agents of anaphylaxis in vaccines: analyzing the formulations of the current authorized COVID-19 vaccines. *J Investig Allergol Clin Immunol*. 2021 Jan 12:0. doi: 10.18176/jiaci.0667.
76. Sellaturay P, Nasser S, Ewan P. Polyethylene Glycol-Induced Systemic Allergic Reactions (Anaphylaxis). *J Allergy Clin Immunol Pract*. 2020 Oct 1:S2213-2198(20)31007-2. doi: 10.1016/j.jaip.2020.09.029.

Legends:

Table 1. Possible predictive risk factors to identify patients with hazard for severe food allergic reactions through nonin gestant routes of exposure
Modified from Liccardi G et al. Severe systemic allergic reaction induced by accidental skin contact with cow milk in a 10-year-old boy. A case-report. *J Investig Allergol Clin Immunol*. 2004; 14: 168-71 (13).

Figure 1. Possible correlation between latex, oxytocin sensitization and airway inflammatory condition including suggested pre-operative diagnostic / therapeutic interventions.
Modified from Liccardi G, Bilò M, Mauro C, Salzillo A, Piccolo A, D'Amato M, Liccardi A, D'Amato G. Oxytocin: an unexpected risk for cardiologic and broncho-obstructive effects, and allergic reactions in susceptible delivering women. *Multidiscip Respir Med*. 2013; 8: 67 (24).

585
586
587
588
589
590
591
592
593
594
595
596

Figure 2. Possible modalities of inducing allergic sensitization through intimate behaviour.
Modified from Liccardi G, Caminati M, Senna G, Calzetta L, Rogliani P. Anaphylaxis and intimate
behaviour. *Curr Opin Allergy Clin Immunol.* 2017; 17: 350-355 (45).

Manuscript accepted for publication