Probiotic as an adjuvant therapy in chronic urticaria: A blinded randomized controlled clinical trial

Najmolsadat Atefi^{1¶}, Morteza Fallahpour^{2¶}, Shokoufeh Sharifi^{1*}, Mohammadreza Ghassemi¹, Masoumeh Roohaninasab¹, Azadeh Goodarzi^{1*}

Affiliation:

1. MD., Department of Dermatology, School of Medicine, Iran University of Medical Sciences (IUMS), Rasool Akram Medical Complex, Tehran, Iran
2. MD., Department of Allergy and Clinical Immunology, School of Medicine, Iran University of Medical Sciences (IUMS), Rasool Akram Medical Complex, Tehran, Iran

¶ First authors:

Najmolsadat Atefi, and Morteza Fallahpour

These authors contributed equally and share first authorship

Corresponding Authors:

*Azadeh Goodarzi

*Shokoufeh Sharifi

-Azadeh Goodarzi*, Department of Dermatology, Rasool Akram Medical Complex, Sattarkhan st., Tehran, Iran

Mob: 00989123882448 Email: goodarzi.az@iums.ac.ir azadeh_goodarzi1984@yahoo.com

Postal code: 1445613131, Phone number: 02166514001

-Shokoufeh Sharifi*, Department of Dermatology, Rasool Akram Medical Complex, Sattarkhan st., Tehran, Iran

Mob: 00989123431408 Email: sharifi_shokoufeh67@gmail.com

Postal code: 1445613131, Phone number: 02166514001

Probiotic as an adjuvant therapy in chronic urticaria: A blinded randomized controlled clinical trial

Abstract:

Background: Chronic spontaneous urticaria (CSU) is a common and treatment challenging disorder which may involve about 2% of normal population and in 50% do not respond properly even to the second line therapies. We aimed to evaluate the efficacy and safety of a symbiotic (prebiotic + probiotic) named as LactoCare in treatment of CSU in the RCT for the first time.

Methods: This blinded RCT conducted on 42 patients (21 patients in control antihistamine group and 21 in intervention antihistamine + probiotic group) with CSU during 8 weeks. The efficacy assessed by Urticaria Activity Score (UAS7) and quality of life measured by Persian validated Dermatology Life Quality Index (DLQI).

Results: Before and after UAS7 score in control group was 35.33 ± 7.81 and 16.86 ± 13.54 , respectively. There was 53% score reduction in control group. Before and after UAS7 score in intervention group was 32 ± 7.84 and 11 ± 11.41 , respectively. There was 66% score reduction in intervention group. Improvement of DLQI in control and intervention group was 44% and 66%, respectively. At the end, UAS7 score reduction and DLQI improvement in both groups was statistically significant.

Conclusion: Probiotics are effective, safe and satisfactory adjuvant therapy for CSU. Combination of probiotic and antihistamines had no statistically significant different efficacy than the antihistamine alone, based on UAS7 score. But Patients with combination therapy may experience higher reduction rate of itch, number of urticaria and total UAS7 score that is clinically of great value and is really practical by itself. Patients with combination therapy experienced more improvement of quality of life (DLQI).

Key words: Chronic Spontaneous Urticaria, Probiotic, Prebiotic, Symbiotic, Antihistamine, RCT, UAS score, DLQI, Urticaria, CSU, Trial

1+2=3

Introduction:

It seems that up to 20% of individuals at some point in their life may be affected by urticaria and angioedema. Episodes lasting for less than 6 week are considered acute, whereas those occurring on most days for more than 6 week are considered as chronic urticarial (CU). The etiology, mechanism, causes and therapeutic options are different in acute and chronic urticarial, therefore the distinction is considered of great importance. Acute urticaria is a self-limited condition, and mast cell activation with an allergen, the main activator. Moreover, foods, drug, insect venom or sting and viral infections are the main causes (1-8).

Chronic urticaria has significant burden and great impact on patient's quality of life and is associated with much psychological comorbidity (9-11).

The most common form of management and treatment system of chronic urticaria is symptomatic therapy. In 5 to 50 percent of patients with chronic urticaria, the first-line treatment (one type of antihistamine regimen) may not result in satisfying disease control. Patients with refractory chronic urticaria require a 4-fold increase in the H1 antagonist dose or continue treatment with Omalizumab, Cyclosporine, or Montelukast if urticaria persists. In patients with chronic spontaneous urticaria (CSU) resistant to four-fold usage dose of antihistamines with higher dose consumption, only 49% of patients reported a decrease or resolution of symptoms, and in 20% of patients, side effects were appeared (12-15).

Studies have been shown that *Lactobacillus salivarius* LS01 and compound of 2 probiotics (*Lactobacillus salivarius* LS01 and *Bifidobacterium breve* BR03) are capable of producing and releasing pro-Th-2 cytokines from Th-1 cells, and help to improve T-helper cells type Th1 / Th2 (17-19).

Probiotics are viable microorganisms that have beneficial effects on the body when consumed in sufficient quantities, and as their great feature, they are safe and secure for the host (20).

Probiotics have been studied with promising results in the treatment of atopic dermatitis (AD), acne, eczema, allergic diseases, skin aging, bacterial and fungal infections, chronic wound healing like diabetic foot ulcers (21-26).

Th-2 cells play a critical role in the pathogenesis of allergic reactions and production of urticarial-related antibodies

The high prevalence of CU and its great effects on the quality of life of patients and their family as well as the partially failure of current treatments led us to design and implement the following study.

Based on our literature review, this is the first blinded RCT to evaluate the efficacy and safety of symbiotic for treatment of chronic urticaria.

Methods and Materials

The present interventional study was a parallel study; analyst blinded randomized controlled clinical trial conducted on 42 patients with chronic urticaria at the allergy and dermatology department of Rasool Akram Medical Complex of Iran University of Medical Sciences, between February and December 2019. Sample size was calculated according to the study of Nettis et al, (16) with regard to $P_1 = 71.1$, $P_2 = 28.9$, $\alpha = 0.05$ and power = 90%, then, 36 patients needed to be enrolled in the study. Fifty two patients were enrolled but 10 of them discontinued the study and finally 42 patients finished (each group=21 patients).

Eligibility criteria

Patients with chronic non-autoimmune non-vasculitic urticaria (at least two days a week more than 6 weeks), aged 18 to 45 years, without any serious co-morbidities (such as malignancies, mental illness, hepatitis, endocrine, rheumatologic or other acute and chronic systemic diseases),

not treated with any drugs other than antihistamines, no history of acute gastrointestinal illnesses such as indigestion and mal-absorption and not taking any corticosteroids for any reason were included in the study.

Patient Recruitment

Informed consent was filled prior to recruitment. Liver, thyroid, kidney function in addition to peripheral cell analysis and autoimmunity screening was done. Assays before allocation is listed as: CBC Diff, BUN/Cr, ALT, AST, ALP, ANA, RF, ESR, CRP, TSH, ANTI TPO, UCA, S/E, stool H. pylori Ag, and in the case of abnormality detection due to probability of systemic disorder, they did not enrolled in the study. The recruitment phase was finished within 5 months. It should be notified that patients were free to exit the study at any time by any reasons such as unwillingness to continue the study or due to side effects especially which did not respond to routine approaches including dose reduction or changing drug type.

Random Sequencing and Allocation

Patients were assigned to control (only antihistamine) and intervention groups (antihistamine+symbiotic) through computerized randomization with 1:1 allocation ratio.

Follow-Up and Evaluation:

Study duration was 8 weeks with 2 visits as the first day visit and final visit after 8 weeks. The safety profile for the probiotic strain over 8 weeks of treatment in our patients was consistent with previous observations in patients treated with probiotic supplement (27). Questionnaire 1 was completed for all eligible participants at the beginning of the study including demographic data of patients (age, sex, occupation, weight, marital status, urticaria number, severity, duration, characteristics and its triggering factors, number of involved days per week, previous antihistamine use and its type). By supervision of the main study investigator, in questionnaire 2 and 3, patients recorded the number and severity of urticaria based on Urticaria Activity Score (UAS7) questionnaire (28) and their quality of life based on Persian validated Dermatology Life Quality Index (DLQI) questionnaire (29), respectively. The number and severity of urticaria also quality of life (questionnaires 2 and 3) were recorded by patients in 8th week visit, again.

Detailed information of questionnaires

Questionnaire 2 filled based on the patient's condition at last week which was designed in two columns, the first column is for itch: 0 (no itching), 1 (slight no annoying itching), 2 (medium annoying itching but does not interfere with daily activities), 3 (severe itching that is annoying and interfere with sleep or daily activities) and the next column is related to the number and severity of urticaria: 0 (no lesion), 1 (less than 20 lesions during 24 hours), 2 (20-50 lesions during 24 hours), 3 (over 50 lesions during 24 hours or a large area of the body involved by large interconnected lesions) (28).

UAS7 score= (Itch score + Urticarial number score) x (number of involved days per week):

Range (0-42)

0=no lesion, 1-6= well controlled, 7-15= mild, 16-27=moderate, 28-42=severe

Based on previous studies, at the end of study, the percentage of UAS7 score reduction calculated and $\leq 10\%$, 11-30%, 31-90% and $>90\%$ score reduction considered as no response, mild response, significant response and complete response, respectively.

DLQI score: 10 item: Range (0-30) (29).

Blinding

There was no blinding of patients or investigator in the study so that only data analyst was blind.

Treatment Regimens

Among these three oral antihistamines (cetirizine 10mg, desloratadine 5mg and fexofenadine 180 mg) two of them was selected and given twice a day (for example cetirizine + desloratadine or desloratadine + fexofenadine...). The selection criteria were based on the patient's preference and appropriate previous clinical response. This treatment strategy was the same for both groups. But in intervention group, in addition to similar antihistamines regimen, patients received twice daily oral probiotic capsules named LactoCare, manufactured by Iranian Bio Fermentation Company.

LactoCare capsule is a synbiotic (probiotic + prebiotic), which contains high amounts of many beneficial and safe bacterial strains (*Lactobacillus rhamnosus*, *Lactobacillus casei*, *Lactobacillus acidophilus*, *Bifidobacterium breve*, *Lactobacillus bulgaricus*, *Bifidobacterium longum*, *Streptococcus thermophilus*) plus fructooligosaccharides as prebiotic.

Laboratory Tests

CBC Diff, BUN/Cr, ALT, AST, ALP, ANA, RF, ESR, CRP, TFT, ANTI TPO, U/A, S/E, stool *H. pylori* Ag, test only at the beginning of the study for all participants.

Statistics and Data Analysis

Descriptive results for values are presented as Mean $\pm$ SD or Percent. Independent T-test and Mann-Whitney test were used to compare the two means. Cochran test was used to investigate the differences between the binary qualitative variables during the measured time. Repeated measures ANOVA and LSD were used to evaluate the quantities over time and between groups. P values less than 0.05 were considered statistically significant. All data were analyzed using SPSS 21 software as intention to treat.

Ethics Committee Approval and Study Registration, and Funding

All stages of this research are committed to the Helsinki Declaration and all patients' information is protected. The ethical code of this study was: IRCTUMS.FMD.REC.1398.129 and the IRCT number was: IRCT20190825044613N1. Costs of this trial were paid by Vice- chancellor for research, Faculty of Medicine, Iran University of Medical Sciences.

Results:

The present study was performed on 42 patients with chronic spontaneous urticaria. (21 patients in intervention synbiotic antihistamine group and 21 patients in control antihistamine group). At first, among demographic data, only the mean WBC count and lymphocyte percentage in the intervention group was significantly lower and higher than the control group, respectively (6800 vs. 8400 $P=0.047$)(46% vs. 37% $P=0.038$) (Table1).

The mean UAS7 score was not statistically different between groups, before therapy. Although after therapy, a statistically significant score reduction was observed in both treatment groups ($P<0.001$) and this score reduction was higher in intervention group, with using the independent t-test, final mean UAS7 score and reduction rate of UAS7, were not significantly different between treatment group ($P>0.05$) (Table 2).

There was not any statistically significant difference between two groups regarding the type of antihistamines regimen and the final UAS7 score ($P>0.05$).

The severity of itch, at first and after 8 weeks of therapy was evaluated and the results showed that at the end of the study the itch severity had a significant decreased in both control and intervention group, ($P=0.047$) and ($P<0.001$), respectively. Itching severity was significantly lower in the intervention group than the control ($P<0.05$), but rate of itching decline during the study was not statistically different between two groups ($P=0.162$).

The urticaria number, at first and after 8 weeks of therapy was evaluated and the results showed that at the end of the study, the urticaria number had a significant decreased in control and

intervention group, ($P = 0.001$) and ($P < 0.001$), respectively. The urticaria number was significantly lower in the intervention group than the control ($P < 0.05$), but the rate of urticaria number decline during the study, was not statistically different between two groups ($P = 0.673$). Based on UAS7 categories, in the control group, firstly 15(71.42%) and 6(28.57%) of the patients had severe and moderate CSU, respectively which at the end of the study changed to 6(28.57%), 2(9.52%), 2(9.52%) and 11(52.38%) as severe, moderate, mild and well-controlled CSU, retrospectively.

Based on UAS7 categories, in intervention group, firstly 19(90.47%) and 2(9.52%) of the patients had severe and moderate CSU, respectively which at the end of study changed to 3(15.78%), 4(19.04%), 7(33.33%) and 7(33.33%) as severe, moderate, mild and well-controlled CSU, retrospectively.

So, in the control group, 3(14%), 3(14%), 11(53 %) and 4(19%) of the patients had no, mild, significant and complete response, respectively and in the intervention group, 2(9%), 12(58%) and 7(33%) of the patients had mild, significant and complete response, respectively. In overall, 19% and 33% of the patients in two groups, respectively, had complete therapeutic response.

At the baseline of the study, there was no significant difference between groups regarding distribution of prescribed antihistamines combination regimen ($P > 0.05$) and by X^2 test it was showed that this regimen did not have a confounding effect on the study results.

In Table 3 you can see the mean final patients' quality of life which increased significantly in both treatment groups and in intervention group, quality of life improvement was significantly higher than the control ($p < 0.05$).

In Figure 1, we showed the main change of UAS7 Score (response) and patients' quality of life, before and after therapy in the control and intervention and group.

We did not find any serious or irreversible side effects among all participants of the study. Some patients had complaint of drowsiness and few patients in symbiotic group experienced mild GI discomfort that was not sustainable.

Discussion

Chronic spontaneous urticaria is a common and treatment challenging disorder which may involve about 2% of normal population and in 50% of cases do not respond properly even to the second line therapies (2, 3, 5, 13-16). Regarding these facts, there are many trends to find and evaluate safe therapeutic options with better or additive response rate. The skin is the largest organ in the body that carries hundreds of microorganisms called the skin microbiota. Colonized bacteria react with toll like receptors on the intestinal epithelial cells and dendritic cells which cause activating and signaling of immune cells including macrophages, NK cells, B cells, T helper cells, cytotoxic T cells and regulatory T cells. Intestinal microbiota imbalance can lead to allergic reaction, thus regulating immune system through intestinal microbiota can affect chronic urticaria (17, 18). There is evidence suggesting that alteration of the composition and/or size of the gut microflora may modulate the IgE response to allergens (19). Probiotics are viable microorganisms that have beneficial effects on the body when consumed in sufficient quantities and their great feature is that they are safe and secure for the host (20). Nowadays, there are many promising data regarding the benefits of microbiota regulation and use of prebiotics, probiotic and synbiotics in various fields of medicine including dermatology (21-26). Whereas modern lifestyles have contributed to changes in the composition of the intestinal microflora, diet supplementation with probiotics may counterbalance the Th-2 activity by promoting Th-1 cytokines production and down regulate IgE production via inhibition of IL-4 and IL-5 production (28-30). In cases of CSU, in which autoreactive IgG antibodies against FcεRI, IgE, or

both or autoreactive IgE antibodies against autoallergens are found, these autoantibodies are causative factors, and IgE, FcεRI, and mast cells are unambiguously at the centre of the pathologic process. For the remaining cases of CSU, IgE, FcεRI, and mast cells are also likely to play essential pathologic roles, although the causative factors have not been identified.

Autoimmune processes might be the primary cause of most cases of CSU. Thus, for those cases with a clear autoimmune cause, the reduction of the IgE by the action of probiotics yields the observed therapeutic efficacy. Even for those cases that involve autoimmune response and autoreactive IgE antibodies subtly, they still involve the central pathologic axis of IgE-FcεRI-mast cells, and probiotics similarly render therapeutic effects (31, 32). However, in recent years, several lines of evidence suggest that some bacterial probiotics can modulate the skin immune system (33).

Since there are many evidences regarding the role of microbiota in pathogenesis and course of the CSU specially gut flora, also little case series or trials focusing on this entity, we designed the first RCT to evaluate the efficacy and safety of a symbiotic (prebiotic+probiotics) named as LactoCare in treatment of CSU.

Specific studies about effects of microbiota and symbiotics in course and treatment of chronic urticaria are really rare, so we discuss about the role of symbiotics in other dermatoses at first and then discuss about urticaria with more detail.

A review study by Notay et al (34), conducted in 2017 entitled "probiotics, prebiotics, and synbiotics for the treatment and prevention of adult dermatological diseases". The results indicated that studies were optimistic about the use of some probiotic and prebiotics strains to improve clinical response in symptomatic AD, also as a treatment for acne. In addition, this review emphasis on further research to evaluate how probiotics and prebiotics could be better used in dermatology.

In a study by Rezazadehet al., (35) from Iran, authors investigated the protective effect of Lactobacillus and Bifidobacterium against chronic urticaria. In this study, stool samples of 20 patients with chronic urticaria were compared with 20 age- and sex-matched healthy controls in terms of Lactobacillus, Bifidobacterium and Bacteroides contents. The results showed that there was no significant difference between the frequencies of these bacteria between groups.

Another study was conducted in 2017 by Nabizadehet al. (36) to investigate the relationship between microbiota composition and chronic urticaria. In this study, 20 patients with chronic urticaria and 20 age-matched healthy controls were selected. The PCR of bacterial DNA genome results showed that the frequency of *A muciniphila*, *C leptum* and *F prausnitzii* in the stool of healthy subjects was significantly higher than in patients with chronic urticaria ($P < 0.001$, $P < 0.01$ and $P < 0.05$, respectively).

Nettis et al. (2016) (37), conducted 8 weeks clinical trial on 38 patients with severe CSU to evaluate efficacy and safety of probiotics. They used UAS7 score as primary outcome measure also assessed the patients' quality of life. Patients received double dose of 3 or more types of antihistamines also received oral probiotics named as Bifiderm twice daily. Patients visited at the beginning, 4 weeks and 8 weeks after study started. Nine patients experienced mild clinical improvement (23.7%), one patient reported significant clinical improvement (2.6%) and one patient completely recovered (2.6%). Twenty seven patients showed no signs of recovery (71.1%). In addition, no adverse effects were reported during the study. In this study we evaluated the clinical efficacy and safety of an intake of a capsule is a synbiotic (probiotic + prebiotic), which contains high amounts of many beneficial and safe bacterial strains (*Lactobacillus rhamnosus*, *Lactobacillus casei*, *Lactobacillus acidophilus*, *Bifidobacterium breve*,

Lactobacillus bulgaricus, Bifidobacterium longum, Streptococcus thermophilus) plus fructooligosaccharides as prebiotic in patients with CSU.

The Nettis et al. study (37) was a single arm before-after trial but our study was a RCT, we compared the results of Nettis's study with the results of the intervention arm of ours that showed 2(9%), 12(58%) and 7(33%) of our patients had mild, significant and complete response, respectively. Our positive therapeutic results were more excellent than the previous study and may be due to better case selection, randomization and the least confounders due to better study design, also all cases of Nettis's study initially had severe urticaria with highest UAS7 score but 90% of our CSU group had severe urticaria with mean UAS7 score as 32. In our study improvement of quality of life was higher than the Nettis's study, logically due to better therapeutic response. The safety was comparable with Nettis's study.

In a review article of Ghaffari et al. (2013) (38), the efficacy of similar therapeutic regimens has been mentioned, as that our study confirmed too. The DLQI was conversely related to urticarial severity that is to somehow similar to our findings that final DLQI was higher in the intervention group which had higher UAS7 score reduction.

Baiardini et al. (2003) (39) found that the quality of life in CSU patients were lower than the respiratory allergies in various social, physical and psychological aspects, that is comparable with our results which showed an improvement of DLQI during the study by decreasing of UAS7 score.

In our study although based on the statistical analysis, there were not any significant difference between control and intervention group regarding UAS7 score (final UAS7 score and the reduction rate during the study), but there were many differences in severity category change and response rate of 2 treatment groups during time that are really important in practical managements of patients. So that symbiotics may make the therapeutic results of urticaria better but for more exact interpretations, we need further studies with higher sample size.

There are many articles in dermatology regarding potential role of microbiota and efficacy of prebiotics, probiotics and synbiotics in many dermatoses such as dermatitis, atopic eczema, acne and etc, that in this study we focused on chronic urticaria and designed the first blinded RCT, in this regard (40-44).

Conclusion:

Probiotics and symbiotics are effective, safe and satisfactory adjuvant therapy for CSU, although combination of probiotic and antihistamine did not have significant efficacy difference compared with the antihistamine alone based on final mean UAS7 score, but patients with combination therapy may experience higher mean reduction rate of UAS7 (although insignificant) and also significantly higher reduction rate of itch and number of urticaria, that is clinically really important and practical. In conclusion, our study suggests that a probiotics administered twice daily for 8 weeks might reduce the symptom scores and quality of life scores in a part of patients with CSU who remained symptomatic despite treatment with H1 antihistamine. The probiotic approach might represent a new well tolerated option in the treatment of CSU.

Limitations and recommendations

Loss to follow ups was one of the major limitation of this study, also our study was not patient blinded as we did not have any placebo. High cost of symbiotic was another limitation of our study. With higher sample size and well-designed study it is probable to observe significant difference between routine and combination therapies (+ probiotic or symbiotics), regarding score reduction.

Acknowledgement:

This study was supported by a research grant from Iran University of Medical Sciences, Tehran, Iran. Also, the authors would like to thank Rasool Akram Medical Complex Clinical Research Development Center (RCRDC) for its technical and editorial assist.

Ethics Committee Approval and Study Registration and Funding: All stages of this research are committed to the Helsinki Declaration and all patients' information is protected. The ethical code of this study was: IR.IUMS.FMD.REC.1398.129 and the IRCT number was: IRCT20190825044612N1. Costs of this trial were paid by Vice chancellor for research, Faculty of Medicine, Iran University of Medical Sciences.

Conflict of Interests:

The authors declare that they have no conflict of interest.

Reference:

1. Maurer M, Weller K, Bindslev-Jensen C, Giménez-Arnau A, Bousquet PJ, Bousquet J, et al: Unmet clinical needs in chronic spontaneous urticaria. *Allergy* 2011; 66: 317-30
2. Henz BM: Physical urticaria. In Henz BM, Zuberbier T, Grabbe J, and Monroe E (eds): *Urticaria. Clinical, diagnostic and therapeutic aspects*. Berlin, Springer 1998. 55-89
3. Cho CB, Shutes SA, Altrich MK, Ardoin SP, Phillips G, Ogeogu PU. Autoantibodies in chronic idiopathic urticaria and nonurticarial systemic autoimmune disorders. *Ann Allergy Asthma Immunol* 2013; 110 (1): 29-33.
4. Pite H, Wedi B, Borrego LM, Kapp A, Raap U. Management of childhood urticaria: current knowledge and practical recommendations. *ActaDermVenereol* 2013; 93 (5): 500-8
5. Confino-Cohen R, Chodick G, Leshno M, et al. Chronic urticaria and autoimmunity: associations found in a large population study. *J Allergy ClinImmunol* 2012; 129 (5): 1307-13.
6. Losol P, Yoo HS, Park HS. Molecular genetic mechanisms of chronic urticaria. *Allergy Asthma Immunol Res* 2014; 6: 13-21.
7. Grattan CE, Sabroe RA, Greaves MW. Chronic urticaria. *J Am AcadDermatol* 2002; 46(5):645-57.
8. Kulthanan K, Jiamton S, Thumpimulvatana N, Pinkaew S. Chronic idiopathic urticaria: prevalence and clinical course. *J Dermatol* 2007;34(5):294-301.
9. Pasaoglu G, Bavbek S, Tugcu H, Abaloğlu O, Misirligil Z. Psychological status of patients with chronic urticaria. *J Dermatol* 2006;33(11):765-71.
10. Atefi N, Rahimi M, Peyghambari S, Ghourchian S. Psychological status in patients with chronic urticarial., *MJIRI* 2011; 25 (4):200- 4.
11. Balp MM, Khalil S, Tian P, Gabriel S, Vietri J, Zuberbier T. Burden of chronic urticaria relative to psoriasis in five European countries. *Eur Acad DermatolVenereol* 2018; 32(2):282-90.
12. Mignon T, van der Eijzen , Harmieke van Os, et al. Effectiveness and safety of antihistamines up to fourfold or higher in treatment of chronic spontaneous urticaria . *Clin Transl Allergy* 2017 ;7(1):4.
13. Nettis E, Colanardi MC, Barra L, Ferrannini A, Vacca A, Tursi A. Levocetirizine in the treatment of chronic idiopathic urticaria: a randomized, double-blind, placebo-controlled study. *Br J Dermatol* 2006; 154(3):533-38.
14. Nettis E, Delle Donne P, Di Leo E, Calogiuri GF, Ferrannini A, Vacca A. Rupatadine for the treatment of urticaria. *Expert Opin Pharmacother.* 2013;14(13):1807-13.
15. Zuberbier T, Bindslev-Jensen C, Canonica W, Grattan CE, Greaves MW, Henz BM, et al. EAACI/GA2LEN/EDF. EAACI/GA2LEN/EDF guideline: management of urticaria. *Allergy* 2006;61:319-21.

16. Nettis E, Colanardi MC, Paradiso MT, Ferrannini A. Desloratadine in combination with montelukast in the treatment of chronic urticaria: a randomized, double-blind, placebo-controlled study. *ClinExp Allergy* 2004; 34(9):1401-1407
17. Drago L, Nicola L, Iemoli E, Banfi G, De Vecchi E. Strain-dependent release of cytokines modulated by *Lactobacillus salivarius* human isolates in an in vitro model. *BMC Res Notes* 2010;3:44-7
18. Gregori S, Goudy KS, Roncarolo MG, The cellular and molecular mechanisms of immunosuppression by human type 1 regulatory T cells, *Front Immunol.* 2012 ;3:30
19. Kirjavainen PV, Apostolou E, Arvola T, Salminen SJ, Gibson GR, Isolauri E. Characterizing the composition of intestinal microflora as a prospective treatment target in infant allergic disease. *Immunol Med Microbiol.* 2001;32:1-7.
20. Hil C ,Guarner F ,Reid G ,et al. Expert consensus document :The international Scientific Association for Probiotic consensus statement on the scope and appropriate use of the term Probiotic. *Nat Rev Gastroenterol Hospital* 2014;11(8):506
21. Fuchs Tarlovsky V, Marquez Barba MF ,Sriram K . Probiotics in dermatologic practice .*Nutr* 2016;32(3):289-95.
22. Zutavern A, Brockow I, Schaaf B, Bolte G, von Berg A ,Dieck U, et al. Timing of solid food introduction in relation to atopic dermatitis and atopic sensitization : result from a prospective birth cohort study .*Pediatrics* 2006 ; 117(2):401-11
23. Vandenplas Y, Huys G, Daube G .probiotics.In : *Textbook of Pediatric Gastroenterology , Hepatology and Nutrition*, Springer, 2016: 515-28
24. Futamura M, Thomas KS, Grindlay DJ, Doney LJ, Morley D, Williams HC. Mapping systematic reviews on atopic eczema – an essential resource for dermatology professionals and researchers. *PLos One* 2013 ;8(3):e58484.
25. Allen SJ, Jordan S, Storey M, Thornton CA, Gravenor MB, Garaiova I, et al. Probiotics in prevention of eczema: a randomized controlled trial. *Arch Dis Child* 2014; 99(11) :1014-9.
26. Mousavi ZE, Mousavi SM, Razavi SH, Farnam-Djomeh Z, Kiani H. Fermentation of pomegranate juice by probiotics lactic acid bacteria .*World MicrobBiotechnol* 2011;27(1):123-8.
27. Costa DJ, Marteau P, Amouyal M et al. Efficacy and safety of the probiotic *Lactobacillus paracasei* LP-33 in allergic rhinitis: a double-blind, randomized, placebo-controlled trial (GA2LEN Study). *Eur J Clin Nutr* 2014;68(5):602-7.
28. Hollis K, Proctor C, McBride D, Balp MM, McLeod L, Hunter S, et al. Comparison of urticaria activity score over 7 (UAS7) values obtained from once – daily and twice-daily versions: results from the ASSURE-CSU study . *Am J Clin Dermatol* 2018; 19(2): 267–74.
29. DLQ Different language July 2019. Available from: <https://www.cardiff.ac.uk>
30. Matszaki T, Chin J. Modulating immune responses with probiotic bacteria. *Immunol Cell Biol* 2000;78:67-73.
31. Jain S. Pathogenesis of chronic urticaria: an overview. *Dermatol Res Pract* 2014;2014:674705.
32. Altman K, Chang C. Pathogenic intracellular and autoimmune mechanisms in urticaria and angioedema. *Clin Rev Allergy Immunol* 2013;45(1):47-62.
33. Guéniche A, Philippe D, Bastien P, Blum S, Buyukpamukcu E, Castiel-Higounenc I. Probiotics for photoprotection. *Dermatoendocrinol* 2009;1:275-9
34. Lacey M, Foolad N, Vaughn AR, Sivamani RK. Probiotics, prebiotics, and synbiotics for the treatment and prevention of adult dermatological diseases. *Am J ClinDermatol* 2017;18(6):721-32.

35. Rezazadeh A, Shahabi S, Bagheri M, Nabizadeh E, Jazani NH. The protective effect of Lactobacillus and Bifidobacterium as the gut microbiota members against chronic urticaria. *Immunopharmacol* 2018 ;59:168-73.
36. Nabizadeh E, Jazani NH, Bagheri M, Shahabi S. Association of altered gut microbiota composition with chronic urticaria. *Annals of Allergy, Asthma Immunol* 2017;119(1): 48-53
37. Nettis E, Di EL, Pastore A, Distaso M, Zaza I, Vacca M, et al. probiotics and refractory chronic spontaneous urticaria. *Ann AllergyClinImmunol* 2016 48(5):182-187.
38. Ghaffari J. Acute chronic urticarial: review of etiology, pathogenesis, manifestation and treatment. *ClinExc* 2013; 1 (1): 83-102.
39. Baiardini I, Giardini A, Pasquali M, Dignetti P, Guerra L, Specchia C, et al. Quality of life and patients' satisfaction in chronic urticaria and respiratory allergy. *Allergy* 2003; 58(7):621-6
40. Goodarzi A, Mozafarpour S, Bodaghabadi M, Mohamadi M. The potential of probiotics for treating acne vulgaris: A review of literature on acne and microbiota. *Dermatol Ther* 2020; 33(3):e13279.
41. Lee J , Seto D, Bielory L ,Meta- analysis of clinical trials of probiotics for prevention and treatment of pediatric dermatitis atopic .*J Allergy ClinImmunol* 2008 ;121(1):116-211.e11
42. Foolad N, Brezinski EA, Chase EP, Armstrong AW. Effect of nutrient supplementation on atopic dermatitis in children: a systematic review of probiotics ,prebiotics,formula and fatty acid .*JAMA Dermatol* 2013 ;30(3): 350-5.
43. Sharma D, Kober MM, Bowe WP. Anti-Aging Effects of Probiotics. *J Drugs Dermatol* 2016; 15(1):9-12.
44. Lukic J, Chen V, Strahinic I, Begovic J, Lev-Tov H, Davis SC, et al. Probiotics or pro-healers: the role of beneficial bacteria in tissue repair. *Wound Repair Regen* 2017; 25(6): 912–22.

Table 1: Comparison between control and intervention group regarding basic demographic characteristics

Variable	Group	Mean	SD	P Value
Age (year)	Control	40.21	13.74	0.552
	Intervention	36.50	10.75	
Weight (kg)	Control	73.31	11.17	0.181
	Intervention	67.37	14.06	
WBC (count)	Control	8376.11	2164.57	0.047
	Intervention	6830.38	2529.42	
Hb	Control	13.09	1.79	0.564
	Intervention	13.56	2.16	
Neutrophil (%)	Control	50.21	12.36	0.418
	Intervention	46.58	14.43	
Lymphocyte (%)	Control	37.15	12.19	0.038
	Intervention	45.94	12.16	
Eosinophil (%)	Control	3.62	2.57	0.483
	Intervention	3.02	2.39	
Urticaria duration (week)	Control	19.81	50.661	0.931
	Intervention	18.76	22.430	
Days involved with urticaria per week (day)	Control	6.43	2.014	0.729
	Intervention	6.24	1.480	
		-	+	P Value
H.pylori (%)	Control	11(66.7)	7(33.3)	0.215
	Intervention	9(42.9)	12(57.1)	
		-	+	P Value
ANA (%)	Control	20(95.2)	1(4.8)	>0.05
	Intervention	20(95.2)	1(4.8)	
		Female	Male	P Value
Gender (%)	Control	15(71.4)	6(28.6)	0.633
	Intervention	15(71.4)	6(28.6)	
Previous therapeutic regimens (%)		steroids	steroids+ antihistamines	P Value
	Control	11(52.4)	10(47.6)	0.197
	Intervention	16(76.2)	5(23.8)	
Types of prescribed antihistamines regimens (%)		Cetirizine+ Fexofenadine	Cetirizine +Desloratadine	P Value
	Control	11(52.4)	10(47.6)	>0.05
	Intervention	11(52.4)	10(47.6)	

Table 2: Mean UAS7 score before and after therapy in both groups and compare the final UAS7 score between the groups

Group		Mean	N	SD	Score reduction (%)	P Value
Control	UAS7 Before	35.33	21	7.81	53%	<0.001
	UAS7 After	16.86	21	13.54		
Symbiotic	UAS7 Before	32.00	21	7.84	66%	<0.001
	UAS7 After	11.00	21	11.41		
Group			N	Mean	SD	P Value
Final UAS7 score in 8 th week	Control		21	16.86	13.54	0.137
	Symbiotic		21	11.00	11.41	

Table 3: Mean DLQI of patients in control and intervention group before and after therapy and comparing the final DLQI score between 2 groups

Group		Mean	N	SD	Improvement of quality of life (%)	P Value
Control	DLQI Before	17.33	21	6.49	44%	<0.001
	DLQI After	9.71	21	9.24		
Symbiotic	DLQI Before	14.47	21	6.40	66%	<0.001
	DLQI After	5.04	21	5.00		
Group		N	Mean	SD	P Value	
Final DLQI in 8 th week	Control	21	9.71	9.24	0.049	
	Symbiotic	21	5.04	5.00		

Figure legend:

Figure 1: The mean change of UAS7 Score (response) and patients' quality of life, before and after therapy in control and intervention and group

Manuscript accepted for publication