





DANIEL LAORDEN<sup>1</sup> , IRENE HERNÁNDEZ<sup>2,3</sup>, JAVIER DOMÍNGUEZ-ORTEGA<sup>2</sup> ,  
DAVID ROMERO<sup>1</sup> , RODOLFO ÁLVAREZ-SALA<sup>1</sup> , SANTIAGO QUIRCE<sup>2</sup> 

# A real-life cohort of mepolizumab treatment in severe eosinophilic asthma

<sup>1</sup>Department of Pneumology, Hospital La Paz, Universidad Autónoma de Madrid, IdiPAZ, and CIBER of Respiratory Diseases, Madrid, Spain

<sup>2</sup>Department of Allergy, Hospital La Paz, IdiPAZ, and CIBER of Respiratory Diseases, Madrid, Spain

<sup>3</sup>Pfizer, Spain

## KEY WORDS

*Asthma; anti-IL5; oral systemic corticosteroids; mepolizumab; eosinophils.*

## Corresponding author

Daniel Laorden  
Department of Pneumology  
Hospital La Paz  
Universidad Autónoma de Madrid, IdiPAZ, and CIBER  
of Respiratory Diseases  
Paseo de la Castellana, 261  
28046 Madrid, Spain  
ORCID: 0000-0001-9054-5925  
E-mail: laordendaniel1@gmail.com

## Doi

10.23822/EurAnnACI.1764-1489.289

## IMPACT STATEMENT

*Our results confirm the efficacy and safety of mepolizumab in routine clinical practice and we observed the significant decrease of ECP, suggesting a reduction in eosinophil activity following mepolizumab treatment.*

## Summary

**Background.** Mepolizumab, a monoclonal antibody that interacts with IL-5, was the first anti-IL-5 approved for uncontrolled severe eosinophilic asthma. In several randomized, placebo-controlled trials, treatment with mepolizumab has shown a significant improvement in asthma symptoms and the need to use of oral corticosteroids (OCS). Several studies have correlated blood levels of eosinophil cationic protein (ECP) with the degree of eosinophilic inflammation, which could make it an indirect marker of eosinophilic activity. **Methods.** This was a single-center retrospective study that included all patients diagnosed with severe eosinophilic asthma under treatment with mepolizumab. We recorded the number of exacerbations, daily prednisone intake, asthma control test scores and forced expiratory volume in the first second. **Results.** We followed 22 patients, 14 of whom were OCS-dependent with a mean daily dose of  $15.85 \pm 15.62$  mg prednisone. After 12 months, only five continued taking OCS and the mean daily dose was reduced by up to  $2.50 \pm 3.84$  mg ( $p < 0.007$ ). The exacerbation rate at baseline was  $2.91 \pm 2.27$  and decreased to  $0.82 \pm 1.14$  in the following year ( $p < 0.001$ ). ACT scores increased significantly from  $16.00 \pm 5.85$  to  $20.71 \pm 4.45$  after six months ( $p = 0.003$ ). We also observed a decrease in ECP from  $81.46 \pm 43.99$   $\mu\text{g/L}$  to  $19.12 \pm 18.80$   $\mu\text{g/L}$  ( $p > 0.001$ ). **Conclusions.** These real-life results are consistent with previous clinical trials demonstrating the efficacy and safety of mepolizumab in routine clinical practice for severe uncontrolled eosinophilic asthma. We observed a significant decrease in blood eosinophil counts and in ECP levels, suggesting a reduction in eosinophil activity following mepolizumab treatment.

## Introduction

Asthma is a common chronic airways disease characterized by variable respiratory symptoms and airflow limitation (1). The worldwide prevalence ranges between 1% and 18%, being highest in developed countries (2). Approximately 4% of this total adult asthma population have a severe asthma phenotype, characterized by difficulty in achieving disease control despite

high-dose inhaled corticosteroids combined with long-acting  $\beta_2$ -agonists (LABAs) or oral corticosteroids (OCS) (3, 4). Such patients may need another type of treatments, such as the new biologic therapies targeting different pathways.

Since omalizumab, the first commercialized biological therapy targeting Immunoglobulin (Ig)E, several antibodies have been developed in the last few years to treat asthma driven by type-2 inflammatory mechanisms, characterized by the production

of type-2 cytokines: interleukin (IL)-4, IL-5 and IL-13. IL-5 plays a key role in stimulating the production, recruitment and activation of eosinophil granulocytes (5). Mepolizumab, a humanized IgG1/k monoclonal antibody that interacts with IL-5 by blocking the binding of this cytokine to the alpha subunit of its receptor (6), was the first IL-5 antibody approved by both the U.S. Food and Drug Administration (FDA) and European Medicines Agency (EMA) for uncontrolled severe eosinophilic asthma. In several randomized placebo-controlled trials, treatment with mepolizumab has shown a significant reduction in exacerbation rates, as well as in the intake of oral corticosteroids and an improvement in the quality of life of these patients (7, 8). Until now, a blood eosinophil count of  $\geq 150/\mu\text{L}$  is practically the only available biomarker for selecting asthma patients who would benefit from treatment with mepolizumab (9). In addition, greater treatment effects with increasing blood eosinophil count have been found. Valid biomarkers in asthma are currently needed to provide information about diagnosis, prognosis, clinical outcomes, and other clinical features, but only a few of them have been described and characterized to date (*e.g.*, IgE, fractional exhaled nitric oxide [FeNO], periostin) (10-12). Several studies have correlated blood levels of eosinophil cationic protein (ECP) to the degree of eosinophilic inflammation, which could make it an indirect marker of eosinophilic activity (12, 13) but this needs further validation in larger populations with severe asthma, its prize is similar to Immunoglobulin E test and ECP could be an additional test to include in a laboratory protocol in an asthma clinic.

Few studies have been published to date about real-life experiences and possible criteria for treatment response with mepolizumab. Our aim was to evaluate severe eosinophilic asthma patients treated with mepolizumab in real-world practice and to find potential predictors of response.

## Materials and methods

Our single-center retrospective study included all patients diagnosed with severe eosinophilic asthma with inadequate control despite treatment with high-dose inhaled corticosteroids (ICS) combined with long-acting beta 2 agonists (LABA) that had started treatment with mepolizumab, according to its current indication in Spain. They were followed up for at least one year in our difficult-to-control asthma unit at the La Paz University Hospital in Madrid, Spain. For inclusion, patients had to have a blood eosinophil count of at least 150 eosinophils/ $\mu\text{L}$  at screening, and/or at least 300 eosinophils/ $\mu\text{L}$  at any point in the previous year. Most of our patients had received continuous or near-continuous treatment with OCS.

Patients received subcutaneous injections of mepolizumab 100 mg every four weeks over a period of at least 12 months. Throughout this treatment period, the following parameters were recorded at baseline, 4-6 months and 12 months: number of asthma exacer-

bations, daily intake (mg) of prednisone, forced expiratory volume in the first second (FEV<sub>1</sub>), blood eosinophil count, ECP levels, total serum IgE levels, FeNO and asthma control test (ACT) scores. At each injection appointment, patients were asked about the occurrence of adverse effects and subsequently registered.

## Statistical analysis

Statistical analysis regarding categorical variables was stated as absolute numbers (n) and percentages (%), and continuous variables as means  $\pm$  standard deviation (SD).

**Table 1** - Baseline demographic and clinical characteristics of severe asthmatics patients.

| Characteristics  | Mepolizumab patients<br>(n = 22) |
|--|----------------------------------|
| Age (years)  | 55.77 $\pm$ 15.03                |
| Female, n (%)  | 15 (68)                          |
| Smoking history, n (%)                                       |                                  |
| Active smoker  | 1 (5)                            |
| Former-smoker  | 10 (45)                          |
| Non-smoker   | 11 (50)                          |
| Body mass index  | 25.39 $\pm$ 5.34                 |
| Continuous OCS therapy, n (%)                                | 14 (64)                          |
| Mean daily dose of prednisone or equivalent at baseline (mg) | 15.85 $\pm$ 15.62                |
| Blood Eosinophils (cells/ml)                                 | 790.91 $\pm$ 549.96              |
| FEV <sub>1</sub> (l) (pre-BD)                                | 1.93 $\pm$ 0.88                  |
| FeNO (ppb)   | 59.99 $\pm$ 63.17                |
| Total IgE (kU/L)   | 214.96 $\pm$ 211.54              |
| ECP ( $\mu\text{g/L}$ )                                      | 81.47 $\pm$ 43.99                |
| ACT  | 16.00 $\pm$ 5.85                 |
| Comorbidities, n (%)   |                                  |
| Rhinitis   | 14 (64)                          |
| Rhinosinusitis   | 17 (77)                          |
| Nasal polyps   | 12 (55)                          |
| Gastroesophageal reflux                                      | 10 (46)                          |
| Bronchiectasis   | 16 (73)                          |
| Obstructive sleep apnea syndrome                             | 7 (32)                           |
| Aspirin-exacerbated respiratory disease (confirmed)          | 2 (9)                            |
| Psychiatric disorder (depression/anxiety)                    | 6 (27)                           |
| Asthma and COPD overlap (ACO)                                | 1 (5)                            |
| Eosinophilic granulomatosis with polyangiitis (confirmed)    | 1 (5)                            |

ACT: asthma control test; BD: bronchodilator; COPD: chronic obstructive pulmonary disease; ECP: eosinophil cationic protein; FeNO: fractional exhaled nitric oxide; FEV1: forced expiratory volume in the first second; IgE: immunoglobulin E; OCS: oral corticosteroids.

The evolution of parameters of interest was calculated using generalized linear models. A P-value < 0.05 was considered statistically significant. Data analysis was performed with SAS 9.3 (SAS Institute, Cary, NC, USA).

### Ethics

The research was conducted ethically in accordance with the World Medical Association Declaration of Helsinki. The protocol was approved by local regulatory ethics committee (PI-2932). Information revealing the patient's identity has been avoided. All patients have been identified by numbers or aliases and not by their real names.

Ethics approval was not required because it was a retrospective and observational study. We did not change our daily clinical practice.

The study participants have given their written informed consent to publish their case.

### Results


We followed 22 patients with severe eosinophilic asthma in our difficult-to-control asthma unit, starting treatment with mepolizumab. Three patients (13.64%) discontinued treatment before completing 12 months. One of them developed herpes zoster after the second injection of mepolizumab, which resolved properly without sequelae. We believed that

this was probably related to the drug. Another patient suffered from non-specific arthralgias that were not considered clearly related to mepolizumab, and the last case involved a non-responder patient after the eighth administration. Other mild side effects, sporadically reported by our patients, included headache, low back pain and local irritation at the injection site.

Demographic and clinical characteristics of all patients, including comorbidities, are shown on **table I**. In this cohort, rhinitis and chronic rhinosinusitis with nasal polyps were the most frequent comorbidities, with a high prevalence of bronchiectasis (73%). Thirteen patients had previously received omalizumab without success.


Prior to mepolizumab treatment, 14/22 patients were OCS-dependent, with a mean daily dose of  $15.85 \pm 15.62$  mg prednisone or equivalent. After 12 months, only 5/22 continued taking OCS and the mean daily dose was reduced by up to  $2.50 \pm 3.84$  mg ( $p < 0.007$ ) (**figure 1**). The exacerbation rate at baseline was  $2.91 \pm 2.27$  and decreased to  $0.82 \pm 1.14$  in the following year ( $p < 0.001$ ) (**figure 2**). The variation of different clinical parameters from baseline to month 6 and 12 is shown in **table II**. Disease control measured by ACT scores significantly increased from  $16.00 \pm 5.85$  to  $20.71 \pm 4.45$  after six months ( $p = 0.003$ ), although a statistically significant change was not obtained at the 12-month follow-up visit. Regarding lung function, FEV<sub>1</sub> showed an average increase of

**Figure 1** - Effect of mepolizumab on OCS intake.



Mean daily prednisone doses (mg) assessed at baseline, six months and after a year of treatment ( $p < 0.007$ ).

**Figure 2** - Exacerbation rate at baseline and after a year of treatment with mepolizumab ( $p < 0.001$ ).



238.85 mL after 12 months of treatment, without reaching statistically significant differences compared to baseline ( $p = 0.261$ ). FeNO measurements increased from a mean of  $59.99 \pm 63.17$  parts per billion (ppb) to  $73.23 \pm 48.28$  ppb after 12 months ( $p = 0.604$ ). All these findings are consistent with a dramatic reduction in the blood eosinophil count after one year, from  $790.91 \pm 549.96$  cells/ml to  $98.00 \pm 90.73$  cells/ml ( $p > 0.001$ ). We also observed a significant decrease in ECP, from  $81.46 \pm 43.99$   $\mu\text{g/L}$  to  $19.12 \pm 18.80$   $\mu\text{g/L}$ , parallel to the reduction in peripheral blood eosinophil count ( $p > 0.001$ ). We observed no significant differences in total IgE at 6 or 12 months.

### Discussion and conclusions

These real-life results are consistent with previous clinical trials showing a clear benefit of this biologic therapy in patients with severe eosinophilic asthma not adequately controlled with high-dose ICS-LABA and/or OCS. A recent article by Bagnasco *et al.* (14) compared the baseline characteristics

of patients treated with mepolizumab in real life with those patients who took part in the main regulatory trials, finding several differences such as improved lung function, higher eosinophil count and treatment with lower doses of prednisone in the former group. This real-world experience is important to support the results of clinical trials and to include new knowledge from clinical practice in guidelines.

The efficacy and safety of mepolizumab have been reported in several clinical trials. In the DREAM study (7), an intravenous formulation resulted in a 48% reduction in clinically significant exacerbations in the active arm *versus* placebo. In fact, high blood eosinophil counts have been found to play an important role in the risk of asthma exacerbations (15, 16); therefore, one of the key therapeutic goals for these types of drugs is to maintain blood eosinophil numbers below relevant thresholds. In our patients, we observed a significant decrease in blood eosinophil counts and in ECP levels, suggesting a reduction in eosinophil activity. However, there is not sufficient evidence to suggest that a decrease in ECP levels is related to a better response to mepolizumab (17). Regarding lung function, the DREAM study confirmed some other early studies in which there was no significant change in FEV1 during treatment (7, 18, 19), as was observed in our patients. More recent clinical trials such as MENSA (8) and MUSCA (20) reported increases of 98 and 120 mL in FEV1 when comparing patients treated with mepolizumab with placebo. In terms of the steroid-sparing effect, the SIRIUS study showed a median percentage reduction of 50% in the glucocorticoid dose in the mepolizumab group, compared to no reduction in the placebo group (21). Similarly, in our study, the number of OCS-dependent patients dropped from 14 to five after one year of treatment. Results from the clinical trials are consistent with those in patients treated with mepolizumab in real life. Rhinitis and nasal polyps were the most frequent comorbidities in our study, as was reported by Bagnasco *et al.* (22), although we observed a higher prevalence of bronchiectasis (73%) in our patients, we think it is due to our Asthma Clinic Protocol where every patient is studied with chest computed tomog-

**Table II** - Variation of different clinical parameters from baseline to month 6 and 12.

|           | Peripheral blood eosinophil count (cells/ml) | ECP ( $\mu\text{g/L}$ ) | Total IgE (kU/L)    | ACT              | FEV <sub>1</sub> (ml) | FeNO (ppb)        |
|-----------|--|-------------------------|---------------------|------------------|-----------------------|-------------------|
| Baseline  | $790.91 \pm 549.96$                          | $81.47 \pm 43.99$       | $214.96 \pm 211.54$ | $16.00 \pm 5.85$ | $1,930.41 \pm 880.29$ | $59.99 \pm 63.17$ |
| 6 months  | $114.00 \pm 99.07$                           | $22.38 \pm 19.08$       | $128.12 \pm 128.77$ | $20.71 \pm 4.45$ | $2,295.63 \pm 634.61$ | $54.95 \pm 43.45$ |
| 12 months | $98.00 \pm 90.73$                            | $19.12 \pm 18.80$       | $160.26 \pm 108.57$ | $18.50 \pm 5.27$ | $2,169.26 \pm 677.75$ | $73.23 \pm 48.28$ |

All data are expressed in mean and standard deviation (SD); ECP; eosinophil cation protein; ACT; asthma control test; FEV1 (ml): exhaled expiratory volume during the first second; FeNO: fractional exhaled nitric oxide.

raphy, however, not all bronchiectasis presented clinical implications in asthma. In 2020, we published a retrospective study (23) performed in patients of our difficult-to-control asthma unit obtaining a similar prevalence of bronchiectasis (60.4%) measured by chest computed tomography. Regarding other comorbidities, one of our patients had a confirmed diagnosis of eosinophilic granulomatosis with polyangiitis (EGPA) that was satisfactorily controlled with mepolizumab 100 mg every four weeks, despite the results of several open-label or placebo-controlled studies where the effectiveness in the induction and maintenance of remission in EGPA was under mepolizumab 300 mg subcutaneously once a month (23-25).

Some real-world studies evaluating the efficacy of mepolizumab, such as Pelaia *et al.* (27), have demonstrated a significant reduction in the annual asthma exacerbation rate (from 3 to 0), with a concomitant reduction in OCS doses (from 6.25 mg/day of prednisone to 0 mg/day), which is similar to our findings. The REDES study (27), a multicenter trial that evaluated mepolizumab efficacy in eosinophilic asthma, found that exacerbation rates decreased by 77.5% and that 50.6% of patients did not suffer any exacerbations during the 12 months of treatment. In addition, 47.8% of the patients were able to discontinue OCS treatment and the mean daily dose was decreased by 59.9%. In turn, the REALITI A study (28) found a reduction in exacerbation rates of 69%, from 4.63 to 1.43 ( $p < 0.001$ ) and the daily OCS dose decreased from 10 mg/day pre-treatment to 5 mg/day after 21 weeks.

A potential cause of discrepancies with these anti-IL5 drugs could be the assessment of treatment response. Global Initiative for Asthma (GINA) Guidelines recommend re-evaluating treatment response to adjust medications according to response (29). Some authors propose the establishment of response criteria for evaluation of treatment. Drick *et al.* (31) recommended an initial follow-up of six months, in order to identify response criteria. In this study patients were divided into two groups: responder and non-responder, having as a primary criterion the improvement of subjective condition (including their subjective asthma-related symptoms, quality of life, number of exacerbations and improvement of physical fitness), as well as lung function and eosinophils in peripheral blood. Some scientific societies have proposed a response-criteria based on previous studies (31).

In real-world settings, only approximately 30% of severe asthma patients are OCS-dependent, with many patients using OCS in short courses of treatment rather than continuously (32). In our study, only one patient discontinued treatment with mepolizumab due to lack of efficacy, and that was after the eighth dose. In this patient, we observed a notable decrease in the peripheral eosinophil count, therefore

this biomarker should not be used in isolation for response assessment. Consistent with previous studies and clinical trials, adverse events were rare and mostly mild in our study, allowing most of the patients to complete treatment for at least a year (33). We measured inflammatory biomarkers such as FeNO and serum ECP, finding different behaviors, but no significant association with clinical improvement was observed. Unlike some previous studies, we did not observe significant improvement in lung function.

Although the small population size is one of the major limitations of this study, our results reinforce those previously reported in regulatory clinical trials and in the few published real-world studies.

In conclusion, the results from our single-center study confirm the efficacy and safety of mepolizumab in routine clinical practice. As reported in previous trials and real-life setting studies, we observed a significant decrease in peripheral blood eosinophilia, asthma exacerbations, OCS intake and an improvement in asthma control, accompanied by the excellent safety profile and tolerability of the drug.

### Fundings

None.

### Contributions

All authors contributed equally to this work.

### Conflict of interests

The authors declare that they have no conflict of interests.

### References

1. Papi A, Brightling C, Pedersen SE, Reddel HK. Asthma. *Lancet*. 2018;391(10122):783-800. doi: 10.1016/S0140-6736(17)33311-1.
2. Bagnasco D, Ferrando M, Bernardi S, Passalacqua G, Canonica GW. The path to personalized medicine in asthma. *Expert Rev Respir Med*. 2016;10(9):957-65. doi: 10.1080/17476348.2016.1205490.
3. Hekking PW, Wener RR, Amelink M, Zwinderman AH, Bouvy ML, Bel EH. The prevalence of severe refractory asthma. *J Allergy Clin Immunol*. 2015;135(4):896-902. doi: 10.1016/j.jaci.2014.08.042.
4. Quirce S, Plaza V, Picado C, Vennera M, Casafont J. Prevalence of uncontrolled severe persistent asthma in pneumology and allergy hospital units in Spain. *J Investig Allergol Clin Immunol*. 2011;21(6):466-71.
5. Israel E, Reddel HK. Severe and Difficult-to-Treat Asthma in Adults. *N Engl J Med*. 2017;377(10):965-76. doi: 10.1056/NEJMra1608969.
6. Gnanakumaran G, Babu KS. Technology evaluation: mepolizumab, GlaxoSmithKline. *Curr Opin Mol Ther*. 2003;5(3):321-5.

7. Pavord ID, Korn S, Howarth P, Bleecker ER, Buhl R, Keene ON, et al. Mepolizumab for severe eosinophilic asthma (DREAM): a multicentre, double-blind, placebo-controlled trial. *Lancet*. 2012;380(9842):651-9. doi:10.1016/S0140-6736(12)60988-X.
8. Ortega HG, Liu MC, Pavord ID, Brusselle GG, FitzGerald JM, Chetta A, et al. Mepolizumab treatment in patients with severe eosinophilic asthma. *N Engl J Med*. 2014;371(13):1198-207. doi: 10.1056/NEJMoa1403290.
9. Albers FC, Licskai C, Chanez P, Bratton DJ, Bradford ES, Yancey SW, et al. Baseline blood eosinophil count as a predictor of treatment response to the licensed dose of mepolizumab in severe eosinophilic asthma. *Respir Med*. 2019;159:105806. doi: 10.1016/j.rmed.2019.105806.
10. Eguiluz-Gracia I, Tay TR, Hew M, Escribese MM, Barber D, O'Hehir RE, et al. Recent developments and highlights in biomarkers in allergic diseases and asthma. *Allergy*. 2018;73(12):2290-305. doi:10.1111/all.13628.
11. Villaseñor A, Rosace D, Obeso D, Pérez-Gordo M, Chivato T, Barbas C, et al. Allergic asthma: an overview of metabolomic strategies leading to the identification of biomarkers in the field. *Clin Exp Allergy*. 2017;47(4):442-56. doi: 10.1111/cea.12902.
12. Malinowski A, Fonseca JA, Jacinto T, Alving K, Janson C. Exhaled nitric oxide levels and blood eosinophil counts independently associate with wheeze and asthma events in National Health and Nutrition Examination Survey subjects. *J Allergy Clin Immunol*. 2013;132(4):821-7.e75. doi: 10.1016/j.jaci.2013.06.007.
13. Carlson M, Håkansson L, Peterson C, Stålenheim G, Venge P. Secretion of granule proteins from eosinophils and neutrophils is increased in asthma. *J Allergy Clin Immunol*. 1991;87(1 Pt 1):27-33. doi: 10.1016/0091-6749(91)90209-7.
14. Bagnasco D, Milanese M, Rolla G, Lombardi C, Bucca C, Hefler E, et al. The North-Western Italian experience with anti IL-5 therapy and comparison with regulatory trials. *World Allergy Organ J*. 2018;11(1):34. doi: 10.1186/s40413-018-0210-7.
15. Price DB, Rigazio A, Campbell JD, Bleecker ER, Corrigan CJ, Thomas M, et al. Blood eosinophil count and prospective annual asthma disease burden: a UK cohort study. *Lancet Respir Med*. 2015;3(11):849-58. doi: 10.1016/S2213-2600(15)00367-7.
16. Zeiger RS, Schatz M, Li Q, Chen W, Khatry DB, Gossage D, et al. High blood eosinophil count is a risk factor for future asthma exacerbations in adult persistent asthma. *J Allergy Clin Immunol Pract*. 2014;2(6):741-50. doi: 10.1016/j.jaip.2014.06.005.
17. Büttner C, Lun A, Splettstoesser T, Kunkel G, Renz H. Monoclonal anti-interleukin-5 treatment suppresses eosinophil but not T-cell functions. *Eur Respir J*. 2003;21(5):799-803. doi: 10.1183/09031936.03.00027302.
18. Leckie MJ, ten Brinke A, Khan J, Diamant Z, O'Connor BJ, Walls CM, et al. Effects of an interleukin-5 blocking monoclonal antibody on eosinophils, airway hyper-responsiveness, and the late asthmatic response. *Lancet*. 2000;356(9248):2144-8. doi: 10.1016/s0140-6736(00)03496-6.
19. Flood-Page P, Swenson C, Faiferman I, Matthews J, Williams M, Brannick L, et al. A study to evaluate safety and efficacy of mepolizumab in patients with moderate persistent asthma. *Am J Respir Crit Care Med*. 2007;176(11):1062-71. doi: 10.1164/rccm.200701-085OC.
20. Chupp GL, Bradford ES, Albers FC, Bratton DJ, Wang-Jairaj J, Nelsen LM, et al. Efficacy of mepolizumab add-on therapy on health-related quality of life and markers of asthma control in severe eosinophilic asthma (MUSCA): a randomised, double-blind, placebo-controlled, parallel-group, multicentre, phase 3b trial. *Lancet Respir Med*. 2017;5(5):390-400. doi: 10.1016/S2213-2600(17)30125-X.
21. Bel EH, Wenzel SE, Thompson PJ, Prazma CM, Keene ON, Yancey SW, et al. Oral glucocorticoid-sparing effect of mepolizumab in eosinophilic asthma. *N Engl J Med*. 2014;371(13):1189-97. doi: 10.1056/NEJMoa1403291.
22. Bagnasco D, Caminati M, Menzella F, Milanese M, Rolla G, Lombardi C, et al. One year of mepolizumab. Efficacy and safety in real-life in Italy. *Pulm Pharmacol Ther*. 2019;58:101836. doi: 10.1016/j.pupt.2019.101836.
23. Zamarron E, Romero D, Fernández-Lahera J, Villasante C, Pinilla I, Barranco P, et al. Should we consider paranasal and chest computed tomography in severe asthma patients? *Respir Med*. 2020;169:106013. doi: 10.1016/j.rmed.2020.106013.
24. Wechsler ME, Akuthota P, Jayne D, Khoury P, Klion A, Langford CA, et al. Mepolizumab or Placebo for Eosinophilic Granulomatosis with Polyangiitis. *N Engl J Med*. 2017;376(20):1921-32. doi: 10.1056/NEJMoa1702079.
25. Steinfeld J, Bradford ES, Brown J, Mallett S, Yancey SW, Akuthota P, et al. Evaluation of clinical benefit from treatment with mepolizumab for patients with eosinophilic granulomatosis with polyangiitis. *J Allergy Clin Immunol*. 2019;143(6):2170-7. doi: 10.1016/j.jaci.2018.11.041.
26. Guillevin L. Vasculitis: Mepolizumab for eosinophilic granulomatosis with polyangiitis. *Nat Rev Rheumatol*. 2017;13(9):518-9. doi: 10.1038/nrrheum.2017.117.
27. Pelaia C, Busceti MT, Solinas S, Terracciano R, Pelaia G. Real-life evaluation of the clinical, functional, and hematological effects of mepolizumab in patients with severe eosinophilic asthma: Results of a single-centre observational study. *Pulm Pharmacol Ther*. 2018;53:1-5. doi: 10.1016/j.pupt.2018.09.006.
28. Domingo Ribas C, Carrillo Díaz T, Blanco Aparicio M, Martínez Moragón E, Banas Conejero D, Sánchez Herrero MG; REDES Study Group. REal world Effectiveness and Safety of Mepolizumab in a Multicentric Spanish Cohort of Asthma Patients Stratified by Eosinophils: The REDES Study. *Drugs*. 2021;81(15):1763-74. doi: 10.1007/s40265-021-01597-9.
29. Harrison T, Canonica GW, Chupp G, Lee J, Schleich F, Welte T, et al. Real-world mepolizumab in the prospective severe asthma REALITI-A study: initial analysis. *Eur Respir J*. 2020;56(4):2000151. doi: 10.1183/13993003.00151-2020.
30. Reddel HK, Bacharier LB, Bateman ED, Brightling CE, Brusselle GG, Buhl R, et al. Global Initiative for Asthma Strategy 2021: executive summary and rationale for key changes. *Eur Respir J*. 2021;59(1):2102730. doi: 10.1183/13993003.02730-2021.
31. Drick N, Seeliger B, Welte T, Fuge J, Suhling H. Anti-IL-5 therapy in patients with severe eosinophilic asthma - clinical efficacy and possible criteria for treatment response. *BMC Pulm Med*. 2018;18(1):119. doi: 10.1186/s12890-018-0689-2.
32. Upham JW, Le Lievre C, Jackson DJ, Masoli M, Wechsler ME, Price DB, et al. Defining a Severe Asthma Super-Responder:

- Findings from a Delphi Process. *J Allergy Clin Immunol Pract.* 2021;9(11):3997-4004. doi: 10.1016/j.jaip.2021.06.041.
33. Sullivan PW, Ghushchyan VH, Globe G, Schatz M. Oral corticosteroid exposure and adverse effects in asthmatic patients. *J Allergy Clin Immunol.* 2018;141(1):110-6.e7. doi: 10.1016/j.jaci.2017.04.009.
34. Khurana S, Brusselle GG, Bel EH, FitzGerald JM, Masoli M, Korn S, et al. Long-term Safety and Clinical Benefit of Mepolizumab in Patients With the Most Severe Eosinophilic Asthma: The COSMEX Study. *Clin Ther.* 2019;41(10):2041-56.e5. doi: 10.1016/j.clinthera.2019.07.007.
35. Coman I, Pola-Bibián B, Barranco P, Vila-Nadal G, Dominguez-Ortega J, Romero D, et al. Bronchiectasis in severe asthma: Clinical features and outcomes. *Ann Allergy Asthma Immunol.* 2018;120(4):409-13. doi: 10.1016/j.anai.2018.02.016.